

# GENERICALLY TRANSITIVE ACTIONS ON MULTIPLE FLAG VARIETIES

ROSTISLAV DEVYATOV

ABSTRACT. Let  $G$  be a semisimple algebraic group whose decomposition into a product of simple components does not contain simple groups of type  $A$ , and  $P \subseteq G$  be a parabolic subgroup. Extending the results of Popov [7], we enumerate all triples  $(G, P, n)$  such that (a) there exists an open  $G$ -orbit on the *multiple flag variety*  $G/P \times G/P \times \dots \times G/P$  ( $n$  factors), (b) the number of  $G$ -orbits on the multiple flag variety is finite.

## INTRODUCTION

Let  $G$  be a semisimple connected algebraic group over an algebraically closed field of characteristic zero, and  $P \subseteq G$  be a parabolic subgroup. One easily checks that the  $G$ -orbits on  $G/P \times G/P$  are in bijection with  $P$ -orbits on  $G/P$ . The Bruhat decomposition of  $G$  implies that the number of  $P$ -orbits on  $G/P$  is finite and that these orbits are enumerated by a subset in the Weyl group  $W$  corresponding to  $G$ . In particular, there is an open  $G$ -orbit on  $G/P \times G/P$ . So we come to the following questions: for which  $G$ ,  $P$  and  $n \geq 3$  is there an open  $G$ -orbit on the *multiple flag variety*  $(G/P)^n := G/P \times G/P \times \dots \times G/P$ ? For which  $G$ ,  $P$  and  $n$  is the number of orbits finite?

Notice that if  $G$  is locally isomorphic to  $G^{(1)} \times \dots \times G^{(k)}$ , where  $G^{(i)}$  are simple, then there exist parabolic subgroups  $P^{(i)} \subseteq G^{(i)}$  such that  $G/P \cong G^{(1)}/P^{(1)} \times \dots \times G^{(k)}/P^{(k)}$ . Hence in the sequel we may assume that  $G$  is simple. Moreover, let  $\pi: \tilde{G} \rightarrow G$  be a simply connected cover. Then  $\pi$  induces a bijection between parabolic subgroups  $P \subseteq G$  and  $\tilde{P} \subseteq \tilde{G}$ , namely  $\tilde{P} = \pi^{-1}(P)$ , and an isomorphism  $\tilde{G}/\tilde{P} \rightarrow G/P$ . Also,  $\tilde{G}/\tilde{P}$  may be considered as  $G$ -variety since  $\text{Ker } \pi$  acts trivially on it. In this sense the isomorphism is  $G$ -equivariant. Therefore we may consider only one simple group of each type.

The classification of multiple flag varieties with an open  $G$ -orbit for maximal subgroups  $P$  was given by Popov in [7]. We need some notation to formulate his result. Fix a maximal torus in  $G$  and an associated simple root system  $\{\alpha_1, \dots, \alpha_l\}$  of the Lie algebra  $\mathfrak{g} = \text{Lie } G$ . We enumerate simple roots as in [3]. Let  $P_i \subset G$  be the maximal parabolic subgroup corresponding to the simple root  $\alpha_i$ .

**Theorem 1.** [7, Theorem 3] *Let  $G$  be a simple algebraic group. The diagonal  $G$ -action on the multiple flag variety  $(G/P_i)^n$  is generically transitive if and only if  $n \leq 2$  or  $(G, n, i)$  is an entry in the following table:*

<i>Type of <math>G</math></i>	$(n, i)$
$A_l$	$n < \frac{(l+1)^2}{i(l+1-i)}$
$B_l, l \geq 3$	$n = 3, i = 1, l$
$C_l, l \geq 2$	$n = 3, i = 1, l$
$D_l, l \geq 4$	$n = 3, i = 1, l - 1, l$
$E_6$	$n = 3, 4, i = 1, 6$
$E_7$	$n = 3, i = 7$

In [7], the following question was posed: for which non-maximal parabolic subgroups  $P \subset G$  is there an open  $G$ -orbit in  $(G/P)^n$ ? We solve this problem for all simple groups except for those of type  $A_l$ .

Denote the intersection  $P_{i_1} \cap \dots \cap P_{i_s}$  by  $P_{i_1, \dots, i_s}$ . It is easy to see that  $P_{i_1, \dots, i_s}$  is a parabolic subgroup and that every parabolic subgroup is conjugated to some  $P_{i_1, \dots, i_s}$ .

**Theorem 2.** *Let  $G$  be a simple algebraic group which is not locally isomorphic to  $SL_{l+1}$ ,  $P \subset G$  be a non-maximal parabolic subgroup and  $n \geq 3$ . Then the diagonal  $G$ -action on the multiple flag variety  $(G/P_i)^n$  is generically transitive if and only if  $n = 3$  and  $(G, P)$  is one of the following pairs:*

Type of $G$	$P$
$D_l, l \geq 5$ is odd	$P_{1, l-1}, P_{1, l}$
$D_l, l \geq 4$ is even	$P_{1, l-1}, P_{1, l}, P_{l-1, l}$

Now let us consider actions with a finite number of orbits. Recall that a  $G$ -variety  $X$  is called *spherical* if a Borel subgroup  $B \subseteq G$  acts on  $X$  with an open orbit. It is well-known that the number of  $B$ -orbits on a spherical variety is finite, see [1], [9]. Equivalently, the number of  $G$ -orbits on  $G/B \times X$  is finite if  $X$  is spherical. Therefore, if  $P \subseteq G$  is a parabolic subgroup and  $X$  is a spherical  $G$ -variety, then the number of  $G$ -orbits on  $G/P \times X$  is finite. The classification of all pairs of parabolic subgroups  $(P, Q)$  such that  $G/P \times G/Q$  is spherical is given in [4] and [8]. According to this classification, if  $(G, P_i)$  is an entry in the table from Theorem 1, then  $G/P_i \times G/P_i$  is spherical and hence the number of  $G$ -orbits on  $G/P_i \times G/P_i \times G/P_i$  is finite. In the last section we prove that the number of  $G$ -orbits on  $(G/P)^n$  is infinite if  $n \geq 4$ . We also check directly that if  $(G, P)$  is an entry in the table from Theorem 2, then the number of  $G$ -orbits on  $G/P \times G/P \times G/P$  is infinite. Thus we come to the following result.

**Theorem 3.** *Let  $G$  be a simple algebraic group,  $P \subset G$  be a parabolic subgroup and  $n \geq 3$ . The following properties are equivalent.*

- (1) *The number of  $G$ -orbits on  $(G/P)^n$  is finite.*
- (2)  *$n = 3$ ,  $P$  is maximal, and there is an open  $G$ -orbit on  $G/P \times G/P \times G/P$ .*
- (3)  *$n = 3$ , and  $G/P \times G/P$  is spherical.*

**Corollary 1.** *Let  $n \geq 3$ . The number of  $G$ -orbits on  $(G/P)^n$  is finite if and only if  $n = 3$  and  $(G, P)$  is one of the pairs listed in the following table:*

Type of $G$	$P$
$A_l$	any maximal
$B_l, l \geq 2$	$P_1, P_l$
$C_l, l \geq 3$	$P_1, P_l$
$D_l, l \geq 4$	$P_1, P_{l-1}, P_l$
$E_6$	$P_1, P_6$
$E_7$	$P_7$

Let us mention a more general result for classical groups. Let  $Q_{(1)}, \dots, Q_{(n)}$  be parabolic subgroups in  $G$ . We call the variety  $G/Q_{(1)} \times \dots \times G/Q_{(n)}$  a *generalized multiple flag variety*. The classification of all generalized multiple flag varieties with a finite number of  $G$ -orbits is given in [5] for  $G = SL_{l+1}$  and in [6] for  $G = Sp_{2l}$ .

Proofs of Theorems 2 and 3 use methods developed in [7]. The results concerning existence of an open orbit in a linear representation space in Section 2 may be of independent interest. In several cases for  $G = SO_{2l}$  the existence of an open orbit on a multiple flag variety is checked directly.

**Acknowledgements.** I am grateful to my scientific advisor Ivan V. Arzhantsev for posing the problem and for paying attention to my work. Also I thank professor Vladimir L. Popov for useful comments and Dmitri A. Timashev for bringing my attention to the articles [4] and [8].

## 1. PRELIMINARIES

Let  $G$  be a connected simple algebraic group over an algebraically closed field  $\mathbb{K}$  of characteristic zero and  $\mathfrak{g} = \text{Lie } G$ . Fix a Borel subgroup  $B \subset G$  and a maximal torus  $T \subset B$ . These data determine a root system  $\Phi$  of  $\mathfrak{g}$ , a positive root subsystem  $\Phi^+$  and a system of simple roots  $\Delta \subseteq \Phi^+$ ,  $\Delta = \{\alpha_1, \dots, \alpha_l\}$ . Choose a corresponding Chevalley basis  $\{x_i, y_i, h_i\}$  of  $\mathfrak{g}$ . We have  $[h, x_i] = \alpha(h)x_i$ ,  $[h, y_i] = -\alpha(h)y_i$  for all  $h \in \mathfrak{t} = \text{Lie } T$  and  $h_i = [x_i, y_i]$ .

Let  $I = \{\alpha_{i_1}, \dots, \alpha_{i_s}\} \subseteq \Delta$  be a subset. The Lie algebra of the parabolic subgroup  $P_I := P_{i_1, \dots, i_s}$  is

$$\mathfrak{p} = \mathfrak{b} \oplus \bigoplus_{\alpha \in \Phi_I} \mathfrak{g}_\alpha,$$

where  $\mathfrak{b} = \text{Lie } B$  and  $\Phi_I \subseteq \Phi^-$  denotes the set of the negative roots such that their decomposition into the sum of simple roots *does not* contain the roots  $\alpha_i$ ,  $i \in I$ . For example,  $P_\Delta = B$  and  $P_\emptyset = G$ . It is known that [2, Theorem 30.1] if a parabolic group  $P$  contains  $B$ , then  $P = P_I$  for some  $I \subseteq \Delta$ . Therefore any parabolic subgroup  $P \subseteq G$  is conjugate to some  $P_I$ . If  $P = P_I$  for some  $I \subseteq \Delta$ , we denote by  $P^-$  the parabolic subgroup whose Lie algebra is

$$\mathfrak{p}^- = \mathfrak{t} \oplus \bigoplus_{\alpha \in -\Phi_I \cup \Phi^-} \mathfrak{g}_\alpha.$$

Denote the weight lattice of  $T$  by  $\mathfrak{X}(T)$ . Let  $\mathfrak{X}^+(T)$  be the subsemigroup of dominant weights with respect to  $B$ . Assume first that  $G$  is simply connected. Then  $\mathfrak{X}^+$  is generated by the fundamental weights  $\pi_1, \dots, \pi_l$ . Given a dominant weight  $\lambda$ , denote the simple  $G$ -module with the highest weight  $\lambda$  by  $V(\lambda)$ . If  $G$  is not simply connected, we may consider a simply connected cover  $p: \tilde{G} \rightarrow G$ , the dominant weight lattice  $\mathfrak{X}^+(p^{-1}(T))$  and the highest weight  $\tilde{G}$ -module  $V(\lambda)$ .

Let  $G$  be a simple group and  $P = P_{i_1, \dots, i_s}$  be a parabolic subgroup. Notice that if there is an open  $G$ -orbit on  $(G/P)^n$ , then there exists an open  $G$ -orbit on  $(G/P_i)^n$  for all  $i \in \{i_1, \dots, i_s\}$ . Indeed, since  $P \subseteq P_i$ , one has the surjective  $G$ -equivariant map  $G/P \rightarrow G/P_i$ ,  $gP \mapsto gP_i$ . It induces the surjective  $G$ -equivariant map  $\varphi: (G/P)^n \rightarrow (G/P_i)^n$ , and the image of an open  $G$ -orbit on  $(G/P)^n$  under  $\varphi$  is an open  $G$ -orbit on  $(G/P_i)^n$ . Similarly, if  $G$  acts on  $(G/P)^n$  with an open orbit and  $m < n$ , then  $G$  acts on  $(G/P)^m$  with an open orbit.

Theorem 1 leaves us very few cases of non-maximal parabolic groups to consider. Namely, if  $n > 3$  and  $G$  is of type  $B_l$ ,  $C_l$  or  $D_l$ , then  $G$  never acts on  $(G/P)^n$  with an open orbit. If  $n = 3$  and  $G$  is of type  $B_l$  or  $C_l$ , it suffices to consider  $P = P_{1,l}$ , and we show that there is no open orbit in this case. If  $n = 3$  and  $G$  is of type  $D_l$ , an open orbit may exist only if  $P = P_I$  where  $I \subseteq \{\alpha_1, \alpha_{l-1}, \alpha_l\}$ . So there are four cases to consider. We reduce the case  $P_{1,l-1}$  to the case  $P_{1,l}$ . If  $G$  is of type  $E_6$ , the only parabolic group we should consider is  $P = P_{1,6}$ . We show that there is no open orbit for  $n = 3$ . If  $G$  is of type  $E_7$ , if there existed an open  $G$ -orbit on  $(G/P)^n$  for  $n \geq 3$ , then the only maximal parabolic subgroup containing  $P$  would be  $P_7$ , but in this case  $P$  should be maximal itself. If  $G$  is of type  $E_8$ ,  $F_4$  or  $G_2$ , an open orbit exists for no maximal parabolic subgroups for  $n \geq 3$ , so there are no cases to consider.

Given a group  $G$  acting on an irreducible variety  $X$  with an open orbit, according to [7] we denote the maximal  $n$  such that there is an open  $G$ -orbit on  $X^n$  by  $\text{gtd}(G : X)$ . If  $G$  acts on  $X^n$  with an open orbit, we say that the action  $G : X$  is *generically  $n$ -transitive*.

We make use of the following fact proved by Popov.

**Proposition 1.** [7, Corollary 1 (ii) of Proposition 2] *Let  $G$  be a simple algebraic group,  $P$  be a parabolic subgroup,  $P^-$  be an opposite parabolic subgroup,  $L = P \cap P^-$  be the corresponding Levi subgroup and  $\mathfrak{u}^-$  be the Lie algebra of the unipotent radical of  $P^-$ . If  $P$  is conjugate to  $P^-$ , then  $\text{gtd}(G : G/P) = 2 + \text{gtd}(L : \mathfrak{u}^-)$ .*

We suppose that the group  $SO_l$  acts in the  $l$ -dimensional space and preserves the bilinear form whose matrix with respect to a standard basis is

$$Q = \begin{pmatrix} & & & & 1 \\ & 0 & & & \\ & & 1 & & \\ & & & 1 & \\ & \dots & & & 0 \\ 1 & & & & \end{pmatrix}.$$

We denote the  $l$ -dimensional projective space by  $\mathbf{P}^l$  and the Grassmannian of  $k$ -dimensional subspaces in  $\mathbb{K}^l$  by  $\text{Gr}(k, l)$ .

## 2. EXISTENCE OF AN OPEN ORBIT

**2.1. Groups of type  $B_l$ .** By Theorem 1, it is sufficient to consider the case  $P = P_{1,l}$ . The Dynkin diagram  $B_l$  has no automorphisms, hence  $P$  is conjugate to  $P^-$ . So we may apply Proposition 1, and it suffices to check that  $\text{gtd}(L : \mathfrak{u}^-) = 0$ , i. e.  $L$  acts on  $\mathfrak{u}^-$  with no open orbit.

Let  $G = SO_{2l+1}$ . Then  $L = \mathbb{K}^* \times GL_{l-1}$  and the  $L$ -module  $\mathfrak{u}^-$  can be decomposed into the direct sum  $V_1 \oplus V_2 \oplus V_3 \oplus V_4 \oplus V_5$ . Here  $V_1$  is a  $GL_{l-1}$ -module  $(\mathbb{K}^{l-1})^*$  dual to the tautological one and its  $\mathbb{K}^*$ -weight is 1,  $V_2$  is a trivial one-dimensional  $GL_{l-1}$ -module of weight 1,  $V_3$  is a tautological  $GL_{l-1}$ -module  $\mathbb{K}^{l-1}$  of weight 0,  $V_4$  is a  $GL_{l-1}$ -module  $\mathbb{K}^{l-1}$  of weight 1,  $V_5$  is a  $GL_{l-1}$ -module  $\Lambda^2 \mathbb{K}^{l-1}$  of weight 0. According to this decomposition, we denote components of a vector  $u \in \mathfrak{u}^-$  by  $u_1, u_2, u_3, u_4, u_5$ .

Notice that there exists a  $GL_{l-1}$ -invariant pairing between  $V_1$  and  $V_3$ . Its  $\mathbb{K}^*$ -weight is 1. Also there exists a  $GL_{l-1}$ -invariant pairing between  $V_1$  and  $V_4$ , whose  $\mathbb{K}^*$ -weight is 2. Therefore the rational function

$$\frac{(u_1, u_3)^2}{(u_1, u_4)}$$

is a non-constant invariant for  $L : \mathfrak{u}^-$ , and the action of  $G$  on  $G/P$  is not generically 3-transitive.

**2.2. Groups of type  $C_l$ .** This case is completely similar to the previous one, and again the only thing we should do is to prove that there is no open  $L$ -orbit on  $\mathfrak{u}^-$ , where  $L = P \cap P^-$  is a Levi subgroup of  $P = P_{1,l}$  and  $\mathfrak{u}^-$  is the Lie algebra of the unipotent radical of  $P^-$ .

Let  $G = Sp_{2l}$ . Then  $L = \mathbb{K}^* \times GL_{l-1}$  and the  $L$ -module  $\mathfrak{u}^-$  can be written as  $V_1 \oplus V_2 \oplus V_3 \oplus V_4$ . Here  $V_1$  is a  $GL_{l-1}$ -module  $(\mathbb{K}^{l-1})^*$  and its  $\mathbb{K}^*$ -weight is 1,  $V_2$  is a  $GL_{l-1}$ -module  $\mathbb{K}^{l-1}$  of weight 1,  $V_3$  is a  $GL_{l-1}$ -module  $S^2 \mathbb{K}^{l-1}$  of weight 0,  $V_4$  is a trivial  $GL_{l-1}$ -module of weight 2. According to this decomposition, we denote components of a vector  $u \in \mathfrak{u}^-$  by  $u_1, u_2, u_3, u_4$ .

We see that there exists a  $GL_{l-1}$ -invariant pairing between  $V_1$  and  $V_2$  with  $\mathbb{K}^*$ -weight is 2. Therefore we have the rational invariant

$$\frac{(u_1, u_2)}{u_4}$$

for  $L : \mathfrak{u}^-$ , and the action of  $G$  on  $G/P$  is not generically 3-transitive.

**2.3. Groups of type  $D_l$ .** This time we should consider the following four cases of parabolic subgroups:  $P = P_{1,l-1}, P_{1,l}, P_{l-1,l}, P_{1,l-1,l}$ . One easily checks that  $P$  and  $P^-$  are conjugate except for the cases  $P = P_{1,l}$ ,  $l$  odd, and  $P = P_{1,l-1}$ ,  $l$  odd.

Let  $G = SO_{2l}$ . There exists a diagram automorphism of  $G$  that interchanges  $\alpha_{l-1}$  and  $\alpha_l$ . It preserves the maximal torus and the Borel subgroup and interchanges  $P_{1,l-1}$  and  $P_{1,l}$ . Therefore, the actions  $G : G/P_{1,l-1}$  and  $G : G/P_{1,l}$  are either generically 3-transitive or not generically 3-transitive simultaneously.

2.3.1.  $P = P_{l-1,l}$ . In this case,  $P$  and  $P^-$  are conjugate, and we have to find  $\text{gtd}(L : \mathfrak{u}^-)$ .

The Levi subgroup  $L$  is isomorphic to  $\mathbb{K}^* \times GL_{l-1}$  and the  $L$ -module  $\mathfrak{u}^-$  is isomorphic to  $V_1 \oplus V_2 \oplus V_3$ , where  $V_1$  is a  $GL_{l-1}$ -module  $\Lambda^2 \mathbb{K}^{l-1}$  and its  $\mathbb{K}^*$ -weight is 0,  $V_2$  is a  $GL_{l-1}$ -module  $\mathbb{K}^{l-1}$  of weight 1,  $V_3$  is a  $GL_{l-1}$ -module  $\mathbb{K}^{l-1}$  of weight  $-1$ . We denote components of a vector  $u \in \mathfrak{u}^-$  by  $u_1, u_2, u_3$ .

Let  $l$  be odd. Then a generic element  $u_1 \in V_1$  gives rise to a non-degenerate skew-symmetric bilinear form on the  $GL_{l-1}$ -module  $(\mathbb{K}^{l-1})^*$ . Furthermore, one can consider the corresponding skew-symmetric form on the tautological  $GL_{l-1}$ -module. This form is obtained by matrix inversion and we denote it by  $u_1^{-1}$ . The following function is a rational  $L$ -invariant:

$$u_1^{-1}(u_2, u_3).$$

Thus the action of  $G$  on  $G/P$  is not generically 3-transitive.

Let  $l$  be even. We prove that there is an open  $L$ -orbit on  $\mathfrak{u}^-$ .

Consider the  $GL_{l-1}$ -module  $V' = V_1 \oplus V_2$ , where  $V_2$  is a  $GL_{l-1}$ -module  $\mathbb{K}^{l-1}$  and  $V_1 = \Lambda^2 \mathbb{K}^{l-1}$ .

Since  $l-1$  is odd, the rank of a generic element  $w \in V_1$  is  $l-2$ . Denote the set of all  $w \in V_1$  such that  $\text{rk } w = l-2$  by  $Z$ . Any element  $w \in Z$  gives rise to a (degenerate) skew-symmetric form on  $V_2^*$ , and  $\dim \text{Ker } w = 1$ . Consider the subspace  $(\text{Ker } w)^\perp \subset V_2$  where all the functions from the kernel vanish. Denote  $V_2 \setminus (\text{Ker } w)^\perp$  by  $X_w$ . Clearly,  $W_1 = \cup_{w \in Z} (w \times X_w)$  is an open  $GL_{l-1}$ -invariant subset of  $V'$ .

Let us prove that  $GL_{l-1}$  acts transitively on  $W_1$ . First, given an element  $u = (u_1, u_2) \in W_1$ , one can apply an element of  $GL_{l-1}$  such that the matrix of the bilinear form  $u_1$  in the corresponding basis is

$$R = \begin{pmatrix} 0 & & & & & & & \\ & 0 & 1 & & & & 0 & \\ & -1 & 0 & & & & & \\ & & & \ddots & & & & \\ & & & & 0 & 1 & & \\ & 0 & & & & -1 & 0 & \end{pmatrix}.$$

The first coordinate of  $u_2$  in the new basis is non-zero since  $u_2 \in X_{u_1}$ . Denote the  $i$ -th coordinate of  $u_2$  by  $(u_2)_i$ . The following element of  $GL_{l-1}$  preserves the bilinear form with matrix  $R$ :

$$\begin{pmatrix} 1 & & 0 & \dots & 0 \\ -(u_2)_3/(u_2)_1 & & & & \\ \vdots & & & I_{l-2} & \\ -(u_2)_{l-1}/(u_2)_1 & & & & \end{pmatrix}.$$

When we apply it to  $u_2$ , all its coordinates will be zero except for the first one.

So any element of  $W_1$  can be transformed by  $GL_{l-1}$ -action to an element of the form  $u_1 = R$ ,  $u_2 = ((u_2)_1, 0, \dots, 0)^T$ , where  $(u_2)_1 \neq 0$ . Clearly, all these elements belong to the same  $GL_{l-1}$ -orbit. Call such an element of  $V'$  *canonical*, i. e. call an element  $(u_1, u_2) \in V'$  *canonical* if  $u_1 = R$ ,  $u_2 = ((u_2)_1, 0, \dots, 0)^T$ , where  $(u_2)_1 \neq 0$ . The stabilizer of  $(u_1, \langle u_2 \rangle)$  consists of direct sums of a non-zero  $1 \times 1$  matrix and a symplectic  $(l-2) \times (l-2)$  matrix. Such an element fixes  $u_2$  as well if and only if the first  $1 \times 1$  matrix is 1.

Now we are ready to consider the  $L$ -action on  $\mathfrak{u}^-$ . Maintain the above notation. Since  $V_2$  and  $V_3$  are isomorphic as  $GL_{l-1}$ -modules, for each  $w \in Z \subset V_1$  we can similarly consider the open subset  $V_3 \setminus (\text{Ker } w)^\perp$ . Denote it by  $Y_w$ . Define the subsets  $W_2 = \cup_{w \in Z} (w \times X_w \times Y_w)$  and  $W = \{u \in W_2 : u_2 \text{ is not a multiple of } u_3\}$ . Let us prove that  $L$  acts on  $W$  transitively.

We may suppose that  $u_1$  and  $u_2$  are canonical in the sense stated above. Applying a diagonal matrix from  $(GL_{l-1})_{u_1, \langle u_2 \rangle}$ , we may assume that  $(u_2)_1(u_3)_1 = 1$  since  $V_2$  and  $V_3$  are both tautological  $GL_{l-1}$ -modules. Since  $u_3$  is not a multiple of  $u_2$ , the vector  $v = ((u_3)_2, (u_3)_3, \dots, (u_3)_{l-1})$  is not zero. Since  $Sp_{l-2}$  acts transitively on  $\mathbb{K}^{l-2} \setminus 0$ , there exists an element  $g \in (GL_{l-1})_{u_1, u_2}$ ,  $g = g_1 \oplus g_2$ ,  $g_1 = 1$ ,  $g_2 \in Sp_{l-2}$  such that  $g_2 v = ((u_3)_1, 0, \dots, 0)^T$ . In other words, we may

suppose that  $u_1$  and  $u_2$  are canonical, the only non-zero coordinates of  $u_3$  are the first one and the second one, they are equal, and  $(u_2)_1(u_3)_1 = 1$ .

Now recall that  $L = GL_{l-1} \times \mathbb{K}^*$ , the  $\mathbb{K}^*$ -weights of  $V_1$ ,  $V_2$  and  $V_3$  are 0, 1 and  $-1$ , respectively. Therefore, after applying a suitable element of  $\mathbb{K}^*$ , we have  $u_1 = S$ ,  $u_2 = (1, 0, \dots, 0)^T$  and  $u_3 = (1, 1, 0, \dots, 0)^T$ , and the  $G$ -action on  $G/P$  is generically 3-transitive.

2.3.2.  $P = P_{1,l}$ . In this case, Proposition 1 applies if and only if  $l$  is even.

Let  $l$  be even. It is sufficient to prove that there is an open  $L$ -orbit on  $\mathfrak{u}^-$ .

Again  $L = \mathbb{K}^* \times GL_{l-1}$ , and the  $L$ -module  $V$  can be decomposed into three summands,  $V = V_1 \oplus V_2 \oplus V_3$ , but this time  $V_1$  is a  $GL_{l-1}$ -module  $\Lambda^2 \mathbb{K}^{l-1}$  and its  $\mathbb{K}^*$ -weight is 0,  $V_2$  is a  $GL_{l-1}$ -module  $\mathbb{K}^{l-1}$  of weight 1,  $V_3$  is a  $GL_{l-1}$ -module  $(\mathbb{K}^{l-1})^*$  of weight 1. We denote components of an element  $u \in \mathfrak{u}^-$  by  $u_1, u_2, u_3$ .

Recall the notation we have introduced for the  $GL_{l-1}$ -module  $V'$ . Also this time denote  $Y_w = V_3 \setminus \text{Ker } w$ . Define  $W_2 = \cup_{w \in Z} (w \times X_w \times Y_w)$  and  $W = \{u \in W_2 : \langle u_2, u_3 \rangle \neq 0\}$ . Here  $\langle \cdot, \cdot \rangle$  denotes the  $GL_{l-1}$ -invariant pairing between  $V_2$  and  $V_3$ . Its  $\mathbb{K}^*$ -weight is 2, but the condition  $\langle u_2, u_3 \rangle \neq 0$  is not affected by  $\mathbb{K}^*$ -action, so  $W$  is  $L$ -invariant. We are going to prove that  $L$  acts transitively on  $W$ .

Again we may suppose that  $u_1 = R$  and the only non-zero coordinate of  $u_2$  is the first one. Notice that  $(u_3)_1 \neq 0$  since  $\langle u_2, u_3 \rangle \neq 0$ . This time  $V_2$  and  $V_3$  are dual  $GL_{l-1}$ -modules, so by applying a suitable element of  $(GL_{l-1})_{u_1, \langle u_2 \rangle}$  the coordinates  $(u_2)_1$  and  $(u_3)_1$  can be made equal.

Consider the vector  $v = ((u_3)_2, (u_3)_3, \dots, (u_3)_{l-1})$ . It cannot be zero since  $u_3 \notin \text{Ker } u_1$ . Since  $Sp_{l-2}$  acts transitively on  $(\mathbb{K}^{l-2})^* \setminus 0$ , there exists an element  $g \in (GL_{l-1})_{u_1, u_2}$ ,  $g = g_1 \oplus g_2$ ,  $g_1 = 1$ ,  $g_2 \in Sp_{l-2}$  such that  $g_2 v = ((u_3)_1, 0, \dots, 0)^T$ . In other words, we may suppose that  $u_1$  and  $u_2$  are canonical, the only non-zero coordinates of  $u_3$  are the first one and the second one, and  $(u_2)_1 = (u_3)_1 = (u_3)_2$ .

This time the  $\mathbb{K}^*$ -weights of  $V_2$  and  $V_3$  are both 1, so with the help of the  $\mathbb{K}^*$ -action we can satisfy the equality  $(u_2)_1 = (u_3)_1 = (u_3)_2 = 1$ . Thus,  $L$  acts transitively on  $W$ , and the  $G$ -action on  $G/P$  is generically 3-transitive.

Let  $l$  be odd. Proposition 1 does not apply, and we have to find  $\text{gtd}(G : G/P)$  directly.

Consider the tautological  $SO_{2l}$ -module  $\mathbb{K}^{2l}$ , and let  $e_1, \dots, e_{2l}$  be its standard basis. Let  $X' \subset \text{Gr}(l, 2l)$  be the set of all isotropic subspaces of dimension  $l$  in  $\mathbb{K}^{2l}$ . One easily checks that  $X'$  is a disjoint union of two  $SO_{2l}$ -orbits, and the group  $O_{2l}$  interchanges them. If two subspaces belong to the same  $SO_{2l}$ -orbit, then their intersection is non-zero.

Denote the orbit  $SO_{2l} \langle e_1, \dots, e_l \rangle \subset X'$  by  $X$ . Then  $X$  is an irreducible subvariety in  $\text{Gr}(l, 2l)$ .

For each  $s \in X$  let  $Y_s \subset \mathbf{P}^{2l-1}$  be the set of all lines contained in  $s$ . Clearly,  $W = \cup_{s \in X} (s \times Y_s)$  is a closed  $G$ -invariant subset in  $\text{Gr}(l, 2l) \times \mathbf{P}^{2l-1}$ . One easily checks that  $G/P = W$ .

Let us prove that there exists an open  $G$ -orbit on  $W \times W \times W$ . We impose some conditions on the point  $(s_1, a_1, s_2, a_2, s_3, a_3) \in W \times W \times W$  and so define an open subset  $Y \subseteq W \times W \times W$ . Then we define a point  $p \in \text{Gr}(l, 2l) \times \mathbf{P}^{2l-1} \times \text{Gr}(l, 2l) \times \mathbf{P}^{2l-1} \times \text{Gr}(l, 2l) \times \mathbf{P}^{2l-1}$  and prove that (a) each point  $y \in Y$  belongs to the same  $G$ -orbit that  $p$  does, and (b)  $p$  belongs to  $Y$ . Condition (b) guarantees that  $Y$  is not empty.

Let  $Y \subseteq W \times W \times W$  be the set of all tuples  $(s_1, a_1, s_2, a_2, s_3, a_3)$  such that:

- (1)  $s_1 \cap s_2 \cap s_3 = 0$ .
- (2)  $s_1 + s_2 + s_3 = \mathbb{K}^{2l}$ .
- (3)  $\dim s_1 \cap s_2 = \dim s_2 \cap s_3 = \dim s_1 \cap s_3 = 1$ .
- (4)  $\dim(a_1 + a_2 + a_3) = 3$ .
- (5) The intersection of the subspaces  $s = (s_1 \cap s_2) + (s_2 \cap s_3) + (s_1 \cap s_3)$  and  $a = a_1 + a_2 + a_3$  is zero.
- (6)  $a_i + s_j + s_k = \mathbb{K}^{2l}$ , where  $i = 1, 2, 3, j \neq i, k \neq i, j < k$ .
- (7) The lines  $a_i$  and  $a_j$  are not orthogonal for all  $i \neq j$ .

Notice that if conditions (1)–(3) hold, the sum of subspaces  $s_1 \cap s_2$ ,  $s_2 \cap s_3$  and  $s_1 \cap s_3$  is direct.

Let us prove that  $G$  acts transitively on  $Y$ . Choose vectors  $f_1, f_2, f_3$  such that  $\langle f_i \rangle = a_i$ , and vectors  $f_4, f_5, f_6$  such that  $\langle f_4 \rangle = s_2 \cap s_3$ ,  $\langle f_5 \rangle = s_1 \cap s_3$ ,  $\langle f_6 \rangle = s_1 \cap s_2$ . The restriction of the bilinear form to the subspace  $S = \langle f_1, \dots, f_6 \rangle$  is defined by the following matrix:

$$\begin{pmatrix} 0 & b_1 & b_2 & b_4 & & \\ b_1 & 0 & b_3 & & b_5 & \\ b_2 & b_3 & 0 & & & b_6 \\ b_4 & & & & & \\ & b_5 & & & & \\ & & b_6 & & & \end{pmatrix}.$$

Conditions (6) and (7) imply that  $b_i \neq 0$  for all  $i$ . Clearly, this matrix is non-degenerate.

The above choice of the vectors  $f_i$  allows to multiply them by scalars. Up to scalar multiplication we may assume that all  $b_i = 1$ .

Notice that a cyclic permutation of  $f_1, f_2, f_3$  and the same permutation of  $f_4, f_5, f_6$  performed simultaneously define a linear operator on  $S$  that preserves the restriction of the bilinear form and whose determinant is 1.

Consider the following basis of  $S$ :  $g_1 = f_1$ ,  $g_2 = f_5$ ,  $g_3 = f_6$ ,  $g_4 = f_3 - f_4 - f_5$ ,  $g_5 = f_2 - f_4$ ,  $g_6 = f_4$ . One checks directly that the matrix of the bilinear form with respect to this basis is  $Q$ . Obviously, there exists a matrix  $M$  such that  $(f_1, \dots, f_6) = (g_1, \dots, g_6)M$  and whose elements do not depend on  $a_i$  and  $s_i$ .

The restriction of the bilinear form to  $S$  is non-degenerate, hence its restriction to  $S^\perp$  is also non-degenerate. Since  $s_i = s_i^\perp$ ,  $\dim(s_i \cap S^\perp) = l - 3$  for all  $i$ .

Thus,  $S^\perp$  is a subspace of even dimension equipped with a non-degenerate symmetric bilinear form. We have three isotropic subspaces of maximal dimension in  $S^\perp$ , and the intersection of any two of them is zero. Let us prove the following lemma.

**Lemma 1.** *Let  $(\cdot, \cdot)$  be a non-degenerate symmetric bilinear form in  $\mathbb{K}^{2k}$ , and  $U_1, U_2, U_3$  be isotropic subspaces of dimension  $k$  with  $U_i \cap U_j = 0$  for  $i \neq j$ . Then there exist matrices  $M_1, M_2, M_3 \in Mat_{2k \times k}$  that do not depend on  $U_i$  and a basis  $e_1, \dots, e_{2k}$  of  $\mathbb{K}^{2k}$  such that: (a) the matrix of the bilinear form is  $Q$  and (b)  $(e_1, \dots, e_{2k})M_i$  is a basis of  $U_i$ .*

*Proof.* Consider the non-degenerate linear map  $A: U_1 \rightarrow U_2$  whose graph is the subspace  $U_3$ . This is possible since  $U_i \cap U_j \neq 0$  for  $i \neq j$ . In terms of the map,  $U_3 = \{v + Av \mid v \in U_1\}$ .

Consider the bilinear form  $(v_1, v_2)_A = (v_1, Av_2)$  on  $U_1$ . Since  $U_3$  is isotropic, we have  $0 = (v_1 + Av_1, v_2 + Av_2) = (v_1, v_2) + (Av_1 + Av_2) + (v_1, Av_2) + (Av_1, v_2) = (v_1, Av_2) + (v_2, Av_1) = (v_1, v_2)_A + (v_2, v_1)_A$  for all  $v_1, v_2 \in U_1$ , hence the form  $(\cdot, \cdot)_A$  is skew-symmetric. Assume that it is degenerate and  $v \in U_1$  belongs to its kernel. Then  $(v_1, v) = (v_1, Av) = 0$  for all  $v_1 \in U_1$ . Since the pairing between trivially intersecting isotropic subspaces  $U_1$  and  $U_2$  of maximal dimension is non-degenerate,  $Av = 0$ . Since  $\text{Ker } A = 0$ ,  $v = 0$  and the form  $(\cdot, \cdot)_A$  is non-degenerate.

Thus, we have a symplectic space  $U_1$  with the skew-symmetric form  $(\cdot, \cdot)_A$ . Hence  $k$  is even. Choose a basis  $\langle q_1, \dots, q_k \rangle$  of  $U_1$  such that the matrix of the skew-symmetric form is

$$\begin{pmatrix} & & & & & & & & 1 \\ & & & & & & & & \\ & 0 & & & & & & & \\ & & & & 1 & & & & \\ & & & -1 & & & & & \\ & & & & & & & & \\ & & & & & & & 0 & \\ -1 & & & & & & & & \end{pmatrix}.$$

The vectors  $q_1, \dots, q_k$  are linearly independent, so let them be the first  $k$  elements of a basis of  $\mathbb{K}^{2l}$ . Define the rest of the basis as follows:  $q_{k+j} = -Aq_j$  if  $j = 1, \dots, k/2$  and  $q_{k+j} = Aq_j$  if  $j = k/2 + 1, \dots, k$ . The matrix of the bilinear form  $(\cdot, \cdot)$  is  $Q$ . The subspaces  $U_i$  have the

following bases:

$$\begin{aligned} U_1 &= \langle q_1, \dots, q_k \rangle \\ U_2 &= \langle q_{k+1}, \dots, q_{2k} \rangle \\ U_3 &= \langle q_1 - q_{k+1}, \dots, q_{k/2} - q_{k+k/2}, q_{k/2+1} + q_{k+k/2+1}, \dots, q_k + q_{2k} \rangle. \end{aligned}$$

This completes the proof of the lemma.  $\square$

Consider the following basis of  $\mathbb{K}^{2l}$ :  $g_1, g_2, g_3, q_1, \dots, q_{l-6}, g_4, g_5, g_6$ , where  $q_i$  are defined above in the proof of the lemma. Notice that the matrix of the bilinear form in this basis is  $Q$ . Define the operator  $B: \mathbb{K}^{2l} \rightarrow \mathbb{K}^{2l}$  that maps this basis to the standard one. We know that the matrix of the bilinear form is  $Q$  in both bases, so  $B \in O_{2l}$ .

Let us check that  $\det B = 1$ . Assume that  $\det B = -1$ . Since  $s_1 = \langle g_1, g_2, g_3, q_1, \dots, q_{l-3} \rangle$ ,  $Bs_1 = \langle e_1, \dots, e_l \rangle \in X$ . Since  $s_1 \in X$ , there exists an operator  $C \in SO_{2l}$  such that  $CBs_1 = s_1$ . Thus,  $CB \in (O_{2l})_{s_1}$ ,  $\det CB = -1$ , and the  $O_{2l}$ -orbit  $X'$  cannot be a union of two distinct  $SO_{2l}$ -orbits, a contradiction.

Bases of the subspaces  $Ba_i$  and  $Bs_i$  can be written in terms of  $e_i$  using matrices that do not depend on  $a_i$  and  $s_i$ . Namely, they are the same matrices that we need to write bases of  $a_i$  and  $s_i$  using  $g_i$  and  $q_i$ , and the latter do not depend on  $a_i$  and  $s_i$ . Denote the 6-tuple  $(Bs_1, Ba_1, Bs_2, Ba_2, Bs_3, Ba_3)$  by  $p$ . It suffices to prove that  $p \in Y$ . Conditions (1)–(7) hold by the construction of  $g_i$  and  $q_i$ , but we should check that  $(Bs_1, Bs_2, Bs_3) \in X \times X \times X$ . It is sufficient to find elements of  $SO_{2l}$  that map  $s_1$  to  $s_2$  and  $s_1$  to  $s_3$ . Since  $s_i = (s_i \cap S) \oplus (s_i \cap S^\perp)$ , we find them as direct sums of elements of  $SO(S)$  and  $SO(S^\perp)$ . The elements of  $SO(S)$  are already found, they are cyclic permutations of  $f_1, f_2, f_3$  and  $f_4, f_5, f_6$ . To interchange  $s_1 \cap S^\perp$  and  $s_2 \cap S^\perp$ , consider the map that permutes all the pairs of vectors  $g_i \leftrightarrow g_{2l+1-i}$ ,  $i = 1, \dots, l-3$ . It is orthogonal and its determinant is 1 since  $l-3$  is even. Finally, the operator with the following matrix in the basis  $g_i$  maps  $s_1 \cap S^\perp$  to  $s_3 \cap S^\perp$ .

$$\begin{pmatrix} I_{l-3} & 0 \\ D & I_{l-3} \end{pmatrix}$$

where

$$D = \begin{pmatrix} -1 & & & & & \\ & \ddots & & & & \\ & & -1 & & & \\ & & & 1 & & \\ & 0 & & & \ddots & \\ & & & & & 1 \end{pmatrix}.$$

Therefore,  $SO_{2l}$  acts transitively on  $Y$ , and  $\text{gtd}(G: G/P) = 3$ .

2.3.3.  $P = P_{1,l-1,l}$ . The subgroups  $P$  and  $P^-$  are conjugate for all  $l$ . It is sufficient to find  $\text{gtd}(L: \mathfrak{u}^-)$ , where  $L = (\mathbb{K}^*)^2 \times GL_{l-2}$  and the  $L$ -module  $\mathfrak{u}^-$  is isomorphic to the direct sum of 7 simple modules that we denote by  $V_1, \dots, V_7$ . Namely,  $V_1$  is a  $GL_{l-1}$ -module  $(\mathbb{K}^{l-2})^*$  and its  $(\mathbb{K}^*)^2$ -weight is  $(1, 0)$ ,  $V_2$  is a trivial  $GL_{l-2}$ -module of weight  $(1, 1)$ ,  $V_3$  is a  $GL_{l-2}$ -module  $\mathbb{K}^{l-2}$  of weight  $(0, 1)$ ,  $V_4$  is a trivial  $GL_{l-2}$ -module of weight  $(1, -1)$ ,  $V_5$  is a  $GL_{l-2}$ -module  $\mathbb{K}^{l-2}$  of weight  $(0, -1)$ ,  $V_6$  is a  $GL_{l-2}$ -module  $\mathbb{K}^{l-2}$  of weight  $(1, 0)$ ,  $V_7$  is a  $GL_{l-2}$ -module  $\Lambda^2 \mathbb{K}^{l-2}$  of weight  $(0, 0)$ . Denote the components of  $u \in \mathfrak{u}^-$  by  $u_1, \dots, u_7$ .

There exists a  $GL_{l-2}$ -invariant pairing between  $V_1$  and  $V_3$  whose  $(\mathbb{K}^*)^2$ -weight is  $(1, 1)$ . The following function is a rational  $L$ -invariant:

$$\frac{(u_1, u_3)}{u_4}.$$

Thus, the  $G$ -action on  $G/P$  is not generically 3-transitive.



**2.4. Groups of type  $E_6$ .** The only parabolic subgroup to consider is  $P = P_{1,6}$ . The set  $\{1, 6\}$  of Dynkin diagram vertices is invariant under all automorphisms of the Dynkin diagram. Hence the Weyl group element of the maximal length interchanges  $P$  and  $P^-$ . We have to find  $\text{gtd}(L : \mathfrak{u}^-)$ .

The Levi subgroup  $L$  is locally isomorphic to  $(\mathbb{K}^*)^2 \times SO_8$ , and the  $L$ -module  $\mathfrak{u}^-$  is isomorphic to  $V_1 \oplus V_2 \oplus V_3$ . Here  $V_1$  is an  $SO_8$ -module with the lowest weight  $-\pi_1$ , i. e. a tautological  $SO_8$ -module,  $V_2$  is an  $SO_8$ -module with the lowest weight  $-\pi_3$ ,  $V_3$  is an  $SO_8$ -module with the lowest weight  $-\pi_4$ . Denote the components of  $u \in \mathfrak{u}^-$  by  $u_1, u_2, u_3$ .

Since  $V_1$  is a tautological  $SO_8$ -module, there exists an  $SO_8$ -invariant symmetric bilinear form on it that we denote by  $(u_1, u_1)$ . There exist diagram automorphisms of  $SO_8$  that transform the tautological  $SO_8$ -module to  $SO_8$ -modules isomorphic to  $V_2$  and  $V_3$ . So there exist an  $SO_8$ -invariant on  $V_2$  that we denote by  $(u_2, u_2)$  and an  $SO_8$ -invariant on  $V_3$  that we denote by  $(u_3, u_3)$ . These bilinear forms are not necessarily  $(\mathbb{K}^*)^2$ -invariant, in general their  $(\mathbb{K}^*)^2$ -weights are three pairs of integers. There is a linear combination of these pairs that is equal to zero. Hence, there exists a non-trivial rational  $L$ -invariant of the form

$$(u_1, u_1)^a (u_2, u_2)^b (u_3, u_3)^c,$$

where  $a, b, c \in \mathbb{Z}$ , and the  $G$ -action on  $G/P \times G/P \times G/P$  is not generically transitive.

### 3. FINITE NUMBER OF ORBITS

**Proposition 2.** *Let  $G$  be a simple algebraic group and  $P$  be a proper parabolic subgroup. If  $n \geq 4$ , the number of  $G$ -orbits on  $(G/P)^n$  is infinite.*

*Proof.* Let  $P = P_{i_1, \dots, i_s}$ . Consider the dominant weight  $\lambda = \pi_{i_1} + \dots + \pi_{i_s}$ . Then  $G/P$  is isomorphic to the projectivization of the orbit of the highest weight vector  $v_\lambda \in V(\lambda)$ . In the sequel we shortly write  $i = i_1$ . It is easy to check that  $y_i^2 v_\lambda = 0$ . Denote the unipotent subgroup  $\exp(\mathfrak{t}y_i)$  by  $U_i$ . We see that  $U_i v_\lambda$  is an affine line not containing zero. The closure of its image in the projectivization  $\mathbf{P}(V(\lambda))$  is a projective line  $\mathbf{P}^1 \subseteq G/P \subseteq \mathbf{P}(V(\lambda))$ . Choose  $n \geq 4$  points  $(x_1, \dots, x_n) \in \mathbf{P}^1 \times \dots \times \mathbf{P}^1 \subseteq G/P \times \dots \times G/P$ . The double ratio of the first four of these points does not change under  $G$ -action. Hence, two  $n$ -tuples with different double ratios cannot belong to the same orbit, and the number of orbits is infinite.  $\square$

Now we prove that in the cases  $P = P_{1,l}$  and  $P = P_{l-1,l}$  the number of orbits on  $G/P \times G/P \times G/P$  is infinite.

We suppose that  $G = SO_{2l}$ . Let  $\mathbb{K}^{2l}$  be the tautological  $SO_{2l}$ -module and let  $e_1, \dots, e_{2l}$  be the standard basis. Let  $X' \subset \text{Gr}(l, 2l)$  be the set of all isotropic subspaces of dimension  $l$  in  $\mathbb{K}^{2l}$ . It is known that  $G/P_l$  is isomorphic to a connected component of  $X'$ . In the sequel we suppose that  $G/P_l = X \subseteq X'$ . For each  $s \in X$  let  $Y_s \subset \mathbf{P}^{2l-1}$  be the set of all lines contained in  $s$ . One easily checks that the closed subset  $Y = \cup_{s \in X} (s \times Y_s) \subset \text{Gr}(l, 2l) \times \mathbf{P}^{2l-1}$  is isomorphic to  $SO_{2l}/P_{1,l}$ .

Similarly, if  $s \in X$ , denote by  $Z_s \subset \text{Gr}(l-1, 2l)$  the set of all subspaces of dimension  $l-1$  in  $s$ . Let  $Z$  be the closed subset  $\cup_{s \in X} (s \times Z_s) \subset \text{Gr}(l, 2l) \times \text{Gr}(l-1, 2l)$ . One easily checks that it is isomorphic to  $SO_{2l}/P_{l-1,l}$ .

First, let  $l = 3$ . Consider the following isotropic subspaces:  $S_1 = \langle e_1, e_2, e_4 \rangle$ ,  $S_2 = \langle e_2, e_3, e_6 \rangle$ ,  $S_3 = \langle e_1, e_3, e_5 \rangle$ . They belong to the same  $SO_6$ -orbit, so we may suppose that  $S_1, S_2, S_3 \in X$ . Choose a line  $T_1 \subset S_1$  such that  $T_1 \subset \langle e_1, e_2 \rangle$ . Also choose lines  $T_2 \subset S_2$  and  $T_3 \subset S_3$  such that  $T_2 \subset \langle e_2, e_3 \rangle$  and  $T_3 \subset \langle e_1, e_3 \rangle$ . Impose one more restriction, namely, the sum  $T_2 + T_3$  should be direct and should not be equal to  $\langle e_1, e_2 \rangle$ . Consider the point  $((S_1, T_1), (S_2, T_2), (S_3, T_3)) \in G/P_{1,l} \times G/P_{1,l} \times G/P_{1,l}$ . There are four subspaces of  $\langle e_1, e_2 \rangle$ :  $\langle e_1 \rangle = S_1 \cap S_3$ ,  $\langle e_2 \rangle = S_2 \cap S_3$ ,  $T_1$  and  $T_4 = (T_2 \oplus T_3) \cap \langle e_1, e_2 \rangle$ . Thus, we have defined four lines in  $\mathbb{K}^6$  in terms of intersections and sums of  $S_i$  and  $T_i$ . If we apply an element  $g \in G$  to these four lines, we will obtain four lines defined in the same way using  $gS_i$  and  $gT_i$  instead of  $S_i$  and  $T_i$ . The double ratio of these four lines in their sum of dimension two is not changed under  $G$ -action. Since  $T_1$  is chosen arbitrarily, this double ratio can be any number and the number of orbits is infinite.

Consider the same subspaces  $S_i$  and  $T_i$  and set  $U_1 = T_1 \oplus \langle e_4 \rangle$ ,  $U_2 = T_2 \oplus \langle e_6 \rangle$  and  $U_3 = T_3 \oplus \langle e_5 \rangle$ . The point  $((S_1, U_1), (S_2, U_2), (S_3, U_3))$  belongs to  $Z \times Z \times Z$ . Note that  $\langle e_1, e_2, e_3 \rangle = (S_1 \cap S_2) \oplus (S_2 \cap S_3) \oplus (S_1 \cap S_3)$  and  $T_i = U_i \cap \langle e_1, e_2, e_3 \rangle$ . Again we have a subspace of dimension two and four lines in it defined in terms of intersections and sums of  $S_i$  and  $U_i$ . The existence of  $SO_6$ -invariant double ratio in this case yields that the number of orbits is infinite.

Let  $l > 3$ . Construct the subspaces  $S_i$ ,  $T_i$  and  $U_i$  as above, using the last three basis vectors instead of  $e_4, e_5, e_6$ . Let  $S'_i = S_i \oplus \langle e_4, \dots, e_l \rangle$  and  $U'_i = U_i \oplus \langle e_4, \dots, e_l \rangle$ . The points  $((S'_1, T_1), (S'_2, T_2), (S'_3, T_3))$  and  $((S'_1, U'_1), (S'_2, U'_2), (S'_3, U'_3))$  belong to  $Y \times Y \times Y$  and  $Z \times Z \times Z$ , respectively. Consider also the subspace  $V = (S'_1 \cap S'_2 \cap S'_3)^\perp$ . The restriction of the bilinear form to this subspace is degenerate, its kernel is  $S'_1 \cap S'_2 \cap S'_3 = \langle e_4, \dots, e_l \rangle$ . The quotient is a space of dimension 6 with a bilinear form. The quotient morphism restricted to  $\langle e_1, e_2, e_3, e_{2l-2}, e_{2l-1}, e_{2l} \rangle$  is an isomorphism, so we have subspaces  $S_i$ ,  $T_i$ ,  $U_i$  in the 6-dimensional space. This is exactly the same situation as we had above for the group  $SO_6$ , and it enables us to define double ratios for the points of  $G/P_{1,l} \times G/P_{1,l} \times G/P_{1,l}$  and  $G/P_{l-1,l} \times G/P_{l-1,l} \times G/P_{l-1,l}$  under consideration. Therefore the number of  $SO_{2l}$ -orbits on these multiple flag varieties is infinite. This finishes the proof of Theorem 3.

#### REFERENCES

- [1] M. Brion, *Quelques propriétés des espaces homogènes sphériques*, Manuscripta Math. **55** (1986), no. 2, 191–198.
- [2] J. Humphreys, *Linear algebraic groups*. GTM **21** Springer-Verlag, New York-Heidelberg, 1975.
- [3] J. Humphreys, *Introduction to Lie Algebras and Representation Theory*. GTM **9**, Springer-Verlag, New York-Berlin, 1978.
- [4] P. Littleman, *On spherical double cones*, J. Algebra **166** (1994), no. 1, 142–157.
- [5] P. Magyar, J. Weynman, A. Zelevinsky, *Multiple flag varieties of finite type*, Adv. Math. **141** (1999), no. 1, 97–118.
- [6] P. Magyar, J. Weynman, A. Zelevinsky, *Symplectic multiple flag varieties of finite type*, J. Algebra **230** (2000), no. 1, 245–265.
- [7] V.L. Popov, *Generically multiple transitive algebraic group actions*, Algebraic groups and homogeneous spaces, 481–523, Tata Inst. Fund. Res. Stud. Math., Tata Inst. Fund. Res., Mumbai, 2007.
- [8] J. Stembridge, *Multiplicity-free products and restrictions of Weyl characters.*, Representation Theory **7** (2003), 404–439.
- [9] E.B. Vinberg, *Complexity of actions of reductive groups*, Funkt. Anal. i Pril. **20** (1986), no. 1, 1–13; English transl.: Funct. Anal. Appl. **20** (1986), no 1, 1–11.

CHAIR OF HIGHER ALGEBRA, FACULTY OF MECHANICS AND MATHEMATICS, LOMONOSOV MOSCOW STATE UNIVERSITY, LENINSKIE GORY 1, GSP-1, MOSCOW, 119991, RUSSIA  
*E-mail address:* deviatov@mccme.ru