

ORIENTED SHEAVES ON DOUBLE MOMENT GRAPHS

ROSTISLAV DEVYATOV, MARTINA LANINI, AND KIRILL ZAINOULLINE

ABSTRACT. In the present paper we extend the theory of sheaves on moment graphs due to Braden-MacPherson and Fiebig to the context of an arbitrary oriented equivariant cohomology h (e.g. to algebraic cobordism). We introduce and investigate structure h -sheaves on double moment graphs to describe equivariant oriented cohomology of products of flag varieties. We show that in the case of a total flag variety X the space of global sections of the double structure h -sheaf also describes endomorphism ring of the equivariant h -motive of X .

1. INTRODUCTION

In [GKM] Goresky, Kottwitz and MacPherson showed that the equivariant cohomology of certain varieties equipped with a torus action is completely encoded in the corresponding *moment graph*. We refer to [GKM] for the precise assumptions which are needed, and we limit ourselves to say that all varieties with a finite number of isolated fixed points and one dimensional orbits, and with no ordinary odd dimension cohomology (and hence flag varieties and Schubert varieties) belong to this class of varieties. Under the appropriate hypotheses, the 1-skeleton of the torus action produces a graph, whose vertices are the fixed points and whose edges are closures of 1-dimensional orbits. Clearly, the torus acts on any 1-dimensional orbit via a character (uniquely defined up to a sign, which corresponds to a chosen orientation of the orbit closure) and the graph obtained by labelling the edges by the corresponding torus character is the moment graph. [GKM] theory allows hence to translate computations in equivariant cohomology into a combinatorial problem.

After [GKM], several applications and generalisations of the results therein have appeared. Hypotheses on the variety have been relaxed [GHZ, GH04, Go14] and moment graph techniques have been applied to other cohomology theories [Br97, Ro03, VV03, Kr12, GR13]. If the T -variety also admits a stratification by T -invariant cells, each of which contains a single T -fixed point, then the inclusion relation of cells induces an orientation of the edges. Braden and MacPherson [BMP] proved that the information contained in this oriented moment graph is sufficient to compute the (equivariant) intersection cohomology of the variety. We should notice that intersection cohomology was also treated in [GKM], but no algorithmic combinatorial description was given. In order to obtain this, Braden and MacPherson introduced the notion of *sheaves on moment graphs*, certain algebro commutative objects which depend on the corresponding graph. The equivariant intersection cohomology could be realised as the space of sections of what has later

2010 *Mathematics Subject Classification.* 14F05, 14F43, 14M15.

Key words and phrases. moment graph, equivariant cohomology, motive, projective homogeneous space.

been called the BMP-sheaf, whose stalks give the local equivariant intersection cohomology. Again, it is possible to apply this theory to flag and Schubert varieties, whose intersection cohomology has a prominent role in representation theory. This fact motivated a series of papers by Fiebig, where he developed and axiomatised a sheaf on moment graph theory, and exploited Braden-MacPherson's construction to attack representation theoretical problems (see, for example, [Fi11]).

Let G be a split semisimple linear algebraic group over a field of characteristic zero. In this paper, we propose a theory of sheaves on moment graphs adapted to any G -equivariant theory \mathfrak{h}_G (see Section 5) constructed in [HM13] from an algebraic oriented cohomology theory \mathfrak{h} of Levine-Morel [LM07]. The sheaves we deal with, which we call *oriented cohomology sheaves* or \mathfrak{h} -sheaves, are constituted by collections of modules and maps of modules over $\mathfrak{h}_G(pt)$, satisfying certain conditions which are imposed by the moment graph data. The previously considered sheaves on moment graphs are hence obtained as the special case in which G is an algebraic torus and \mathfrak{h}_G is the equivariant Chow theory. We then use our theory to describe equivariant oriented cohomology of projective homogeneous varieties G/P and their products $G/Q \times G/P$ (here G acts diagonally on the product and P, Q are parabolic subgroups of G) as global sections of certain \mathfrak{h} -sheaves; we also interpret the space of global sections of such sheaves as a *Hom*-space in certain equivariant motivic category.

More precisely, for the homogeneous variety G/P we deal with the *parabolic moment graph* \mathcal{G}^P (see Example 2.3) and its *structure \mathfrak{h} -sheaves* \mathcal{O}^P (see Example 5.4). Observe that the parabolic moment graph is one of the basic and classical tools to describe the T -equivariant cohomology (Chow groups) or K -theory of G/P . As for the products, we introduce a *double moment graph* ${}^Q\mathcal{G}^P$ (see Definition 2.9), and its associated structure \mathfrak{h} -sheaf ${}^Q\mathcal{O}^P$ (Example 5.5). The notion of a double moment graph has not been known before. It turns out to be a useful tool to describe oriented cohomology of the products. Indeed, one of our main results (Theorem 9.1.(B)) says that the space of global sections of the sheaf ${}^Q\mathcal{O}^P$ describes W -invariants (W is the Weyl group of G) of the T -equivariant oriented cohomology of the product.

Following Manin one can introduce the category of G -equivariant \mathfrak{h} -motives (see [PS16, NPSZ] for more details). Let \mathfrak{M}_G be its full additive subcategory generated by motives of G/P for all standard parabolic subgroups P of G . The morphisms in \mathfrak{M}_G are given by cycles in $\mathfrak{h}_G(G/Q \times G/P)$ and their composition is given by the celebrated correspondence product. Our main result (Theorem 9.1) then implies that global sections of the structure sheaves \mathcal{O}^P and ${}^Q\mathcal{O}^P$ describe various *Hom*-spaces between equivariant motives $[G/Q]$ and $[G/P]$ (see Section 10). Moreover, it enables us to interpret the correspondence product in the category \mathfrak{M}_G in moment graphs terms (Proposition 10.7) and to realise the endomorphism ring of the equivariant motive of the total flag variety of type A as the space of global sections of a structure \mathfrak{h} -sheaf on the disjoint union of double moment graphs ${}^Q\mathcal{G}^P$, where Q and P vary on all possible parabolic subgroups of G containing a fixed Borel B (Corollary 10.5).

Acknowledgements. This project started during the workshop Beyond Toric Geometry that was held in May 2017 in Oaxaca, Mexico. M.L. and K.Z. are hence very grateful to the organizers of this workshop and to BIRS Oaxaca for hospitality.

R.D. was partially supported by the Fields Institute for Research in Mathematical Sciences, Toronto, Canada. R.D. and K.Z. were partially supported by the NSERC Discovery grant RGPIN-2015-04469, Canada.

2. DOUBLE MOMENT GRAPHS

Consider a root system Σ with a set of simple roots $\Theta = \{\alpha_1, \dots, \alpha_n\}$ and a set of positive roots Σ^+ . Let W be the Coxeter group generated by simple reflections s_{α_i} . Given two subsets $\Theta_P, \Theta_Q \subset \Theta$ we denote by Σ_P, Σ_Q the respective root subsystems and by W_P and W_Q the respective Coxeter subgroups. We denote by ${}^QW^P$ the set of minimal length double coset representatives of $W_Q \backslash W / W_P$. The Bruhat order ' \leq ' on W descends to ${}^QW^P$ which we can hence consider as a poset itself. Given $u \in {}^QW^P$ consider the subset $\Theta_u = \Theta_Q \cap u(\Sigma_P^+)$ and the respective Coxeter subgroup $W_u = W_Q \cap uW_Pu^{-1}$. Denote by W_Q^u the set of minimal length coset representatives of W_Q/W_u .

We will extensively use the following fact (Double Parabolic Decomposition, see, for example, [Cu85]):

An element y of a double coset $W_Q \backslash W / W_P$ can be uniquely written as $y = wuv$, where $u \in {}^QW^P$, $w \in W_Q^u$ and $v \in W_P$.

In particular, any element of W^P can be uniquely written as wu , where $u \in {}^QW^P$ and $w \in W_Q^u$.

Given $w \in W$ we denote by \bar{w} (resp. \hat{w}) the minimal length coset representative of w in W^P (resp. in ${}^QW^P$).

The following definition has appeared by now in several papers, see, for example, [BMP, Fi08].

Definition 2.1. The data $\mathcal{G} = ((V, \leq), l: E \rightarrow \Sigma^+)$ is called a moment graph if

- (MG1) V is a set of vertices together with a partial order ' \leq ', i.e., we are given a poset (V, \leq) .
- (MG2) E is a set of directed edges labelled by positive roots Σ^+ via the label function l , i.e., an edge $x \rightarrow y \in E$ is labelled by $l(x \rightarrow y) \in \Sigma^+$.
- (MG3) For any edge $x \rightarrow y \in E$, we have $x \leq y$, i.e., direction of edges respects the partial order.

By a moment subgraph of \mathcal{G} we call a moment graph with $(V', \leq) \subset (V, \leq)$, $E' \subset E$ and $l': E' \rightarrow \Sigma^+$ being a restriction of l .

Observe that (MG2) assumes that each edge has a unique label, i.e., there are no multiple edges, and (MG3) implies that \mathcal{G} has no directed cycles.

Example 2.2. Consider the usual Bruhat poset (W, \leq) . The data

$$V := W, E := \{w \rightarrow s_\alpha w \mid w \leq s_\alpha w, \alpha \in \Sigma^+\} \text{ and } l(w \rightarrow s_\alpha w) := \alpha$$

define a moment graph. Observe that the transitive closure of E gives the Bruhat order ' \leq ' and E contains all cover relations of the Bruhat order.

Example 2.3. Given a subset Θ_P of the set of simple roots Θ , consider the Bruhat subposet (W^P, \leq) . The data

$$V = W^P, E = \{w \rightarrow \overline{s_\alpha w} \mid w \leq s_\alpha w, \alpha \in \Sigma^+\} \text{ and } l(w \rightarrow \overline{s_\alpha w}) = \alpha$$

define a moment graph called a parabolic moment graph and denoted by \mathcal{G}^P . Again the transitive closure of E gives the Bruhat order ' \leq ' on W^P and E contains all cover relations of this order.

To show that \mathcal{G}^P has a well-defined label function (i.e. (MG2) holds), let $\Sigma_{\mathbb{R}}$ be a real vector space with an inner product (\cdot, \cdot) that contains Σ . Choose a dominant $\theta \in \Sigma_{\mathbb{R}}$ such that W_P is a stabilizer group of θ . The poset (W^P, \leq) can be identified with the poset $(W\theta, \leq)$, via $w \mapsto w\theta$ (see [St05, Prop. 1.1]), where the partial order on the orbit $W\theta$ is defined by taking the transitive closure of relations

$$\mu \leq s_{\alpha}(\mu) \text{ for all } \alpha \in \Sigma^+ \text{ such that } (\mu, \alpha) > 0.$$

Now suppose that $\overline{s_{\alpha}w} = \overline{s_{\beta}w}$ for $\alpha, \beta \in \Sigma^+$, $w \in W^P$. Then $s_{\alpha}(\mu) = s_{\beta}(\mu)$, where $\mu = w\theta$, which implies that $\alpha = \beta$. In other words, the label α is uniquely determined by the representative $\overline{s_{\alpha}w}$.

Example 2.4. Consider the symmetric group $W = S_3$ generated by simple reflections s_1, s_2 corresponding to simple roots α_1, α_2 . Let $W_Q = W_P = \langle s_1 \rangle$. Then we have ${}^QW^P = \{e, s_2\}$ and

$$\widehat{s_2 e} = \widehat{s_{\alpha_1 + \alpha_2} e} = s_2.$$

Hence, an edge $e \rightarrow s_2$ in the directed graph given by

$$V = {}^QW^P \text{ and } E = \{u \rightarrow \widehat{s_{\alpha}u} \mid u \leq s_{\alpha}u, \alpha \in \Sigma^+\}$$

has to be labelled by two different positive roots α_2 and $\alpha_1 + \alpha_2$, i.e., the label function l of (MG2) is not well-defined. The latter shows that Example 2.3 can not be directly generalized to double cosets.

Indeed, we will show that in certain cases Example 2.3 can be generalized to double cosets if we allow to label the edges $u \rightarrow \widehat{s_{\alpha}u}$ up to an action by the stabilizer group W_u .

First, we introduce the notion of equivariant moment graphs.

Definition 2.5. Given a moment graph $\mathcal{G} = ((V, \leq), l: E \rightarrow \Sigma)$, suppose that a group W acts on the set V and it also acts by permutations on Σ . We say that \mathcal{G} is W -equivariant if

(MGE) For all $w \in W$ and $x \rightarrow y \in E$ if $w(l(x \rightarrow y)) \in \Sigma^+$, then $w(x) \rightarrow w(y) \in E$ and $l(w(x) \rightarrow w(y)) = w(l(x \rightarrow y))$, otherwise $w(y) \rightarrow w(x) \in E$ and $l(w(y) \rightarrow w(x)) = -w(l(x \rightarrow y))$.

Note that a W -equivariant moment graph is not a W -moment graph in the sense of [F15, Def. 4.2].

Lemma 2.6. *The parabolic moment graph \mathcal{G}^P of Example 2.3 is W -equivariant, where W acts on W^P by $w(v) = \overline{wv}$, $v \in W^P$.*

Proof. Suppose $y = \overline{s_{\alpha}x}$ with $x \leq y$ and $\alpha \in \Sigma^+$, then $y = s_{\alpha}xz$, $z \in W_P$ and we have

$$w(y) = w(s_{\alpha}xz) = \overline{s_{w(\alpha)}w(x)}.$$

Since $x \leq y$, we have $x(\theta) = \mu \leq s_{\alpha}(\mu) = y(\theta)$ in $W\theta$, i.e., $(\mu, \alpha) > 0$. Then $(\mu, \alpha) = (w(\mu), w(\alpha)) > 0$ which implies that $w(x)(\theta) = w(\mu) \leq s_{w(\alpha)}w(\mu) = w(y)(\theta)$ if $w(\alpha) \in \Sigma^+$ and $w(\mu) \geq s_{w(\alpha)}w(\mu)$ if $w(\alpha) \in \Sigma^-$. \square

Corollary 2.7. *Let Θ_Q be a subset of Θ and let W_Q be the respective Weyl group. Then the parabolic moment graph \mathcal{G}^P is W_Q -equivariant.*

Second, we define the notion of a closure (an orbit) of a subgraph.

Definition 2.8. Let $\mathcal{G}' = ((V', \leq), l: E' \rightarrow \Sigma^+)$ be a subgraph of a W -equivariant moment graph \mathcal{G} . By a W -closure of \mathcal{G}' we call a subgraph denoted by $W(\mathcal{G}')$ with vertices $\{wx \mid w \in W, x \in V'\}$ and

- edges $w(x) \rightarrow w(y)$ labelled by $w(l(x \rightarrow y))$ for all $x \rightarrow y \in E'$ with $w(l(x \rightarrow y)) \in \Sigma^+$, and
- edges $w(y) \rightarrow w(x)$ labelled by $-w(l(x \rightarrow y))$ for all $x \rightarrow y \in E'$ with $w(l(x \rightarrow y)) \in \Sigma^-$.

Roughly speaking, $W(\mathcal{G}')$ is the smallest W -equivariant moment subgraph containing \mathcal{G}' .

We are now ready to introduce a key object of the present paper

Definition 2.9. Given subsets Θ_P, Θ_Q of the set of simple roots Θ , the data $(({}^QW^P, \leq), l: {}^QE^P \rightarrow \Sigma^+)$ is called the double moment graph and denoted by ${}^Q\mathcal{G}^P$ if it is a moment graph on the set of minimal length double coset representatives with respect to the induced Bruhat order such that the label function l satisfies

$$u' = s_{l(\widehat{u \rightarrow u'})}u, \quad \text{for all } u \rightarrow u' \in {}^QE^P.$$

By definition a double moment graph corresponds to a subgraph \mathcal{G}' of \mathcal{G}^P by

$$E' = \{u \rightarrow \overline{s_{l(\widehat{u \rightarrow u'})}u} \mid u \rightarrow u' \in {}^QE^P\}.$$

Its W_Q -closure $W_Q(\mathcal{G}')$ is called a closure of ${}^Q\mathcal{G}^P$ and denoted by $\langle {}^Q\mathcal{G}^P \rangle$.

To compute the closure of a double moment graph we will use the following

Lemma 2.10. *All edges in a parabolic moment graph \mathcal{G}^P can be obtained by applying W_Q to the edges of the form*

$$u \rightarrow \overline{s_\beta u}, \quad u \in {}^QW^P, \beta \in \Sigma^+.$$

Proof. Consider the W_Q -equivariant moment graph \mathcal{G}^P . Let $\theta \in \Sigma_{\mathbb{R}}$ be a dominant vector with stabilizer W_P . Following [St75, §1B] consider the subposet of $W\theta$

$$(W\theta)_Q = \{\mu \in W\theta \mid (\mu, \alpha) \geq 0 \text{ for all } \alpha \in \Sigma_Q^+\}$$

By [St75, Prop. 1.5] it is isomorphic to the Bruhat subposet $({}^QW^P, \leq)$ via the evaluation map $w \mapsto w\theta$.

Consider an edge $x \rightarrow y$ in \mathcal{G}^P labelled by $\alpha \in \Sigma^+$. Suppose $\widehat{x} \leq \widehat{y}$. By parabolic decomposition, we have $x = wu$, where $w \in W_Q^u$ and $u \in {}^QW^P$. Hence, it corresponds to a vector $w(\mu) \in W\theta$, where $\mu = u(\theta) \in (W\theta)_Q$. Similarly, $y = \overline{s_\alpha x}$ corresponds to a vector $s_\alpha w(\mu)$ in $W\theta$. Since x and y are connected by an edge labelled by α we have $w(\mu) \leq s_\alpha w(\mu)$ that is $(w(\mu), \alpha) = (\mu, w^{-1}(\alpha)) > 0$.

Suppose $w^{-1}(\alpha) \in \Sigma^+$, then the edge $x \rightarrow y$ is obtained from the edge

$$u = w^{-1}x \rightarrow \overline{w^{-1}y} = \overline{s_{w^{-1}(\alpha)}u}$$

labelled by $w^{-1}(\alpha)$ by applying $w \in W_Q^u$ (the latter edge corresponds to $\mu \leq s_{w^{-1}(\alpha)}\mu$ in $W\theta$). If $w^{-1}(\alpha) \in \Sigma^-$, then $\mu \geq s_{w^{-1}(\alpha)}\mu$ which contradicts to the assumption $\widehat{x} \leq \widehat{y}$. \square

3. EXAMPLES OF DOUBLE MOMENT GRAPHS

In the present section we consider parabolic and double moment graphs for root systems of types A and D . For each type we provide necessary and sufficient conditions for $\langle {}^Q\mathcal{G}^P \rangle = \mathcal{G}^P$. We follow the notation of the previous section.

Type A. Let $\Sigma_{\mathbb{R}}$ be a subspace of \mathbb{R}^{n+1} for which the coordinates sum to zero. The Weyl group $W = S_{n+1}$ acts on \mathbb{R}^{n+1} by permutations of coordinates. We fix a set of simple roots $\Theta = \{\alpha_1, \dots, \alpha_n\}$.

Consider the vector $\mu = u(\theta) \in W\theta$, where θ is dominant with the stabilizer W_P and $u \in {}^QW^P$. The subgroup W_Q acts on a vector $\mu = (x_1, \dots, x_{n+1})$ by permutations of coordinates in each of the tuples $\tau_0 = (x_1, \dots, x_{k_1})$, $\tau_1 = (x_{k_1+1}, \dots, x_{k_2})$, \dots , $\tau_r = (x_{k_r+1}, \dots, x_{n+1})$, where $\Theta \setminus \Theta_Q = \{\alpha_{k_1}, \dots, \alpha_{k_r}\}$.

Suppose $\widehat{s_\alpha u} = \widehat{s_\beta u} > u$ for some $\alpha, \beta \in \Sigma^+$. Let $s_\alpha = (i'j')$, $i' < j'$. Since $s_\alpha(\mu) \neq \mu$, we have $x_{i'} \neq x_{j'}$. Since $\widehat{s_\alpha u} \neq u$, the coordinates $x_{i'}$ and $x_{j'}$ belong to different tuples, respectively. Similarly, if $s_\beta = (i''j'')$, $i'' < j''$, then the coordinates $x_{i''} \neq x_{j''}$ belong to different tuples. Since $\widehat{s_\alpha u} = \widehat{s_\beta u}$ and the action by W_Q preserves the numbers of equal coordinates in each tuple, the coordinates $x_{i'}$, $x_{i''}$ (resp. $x_{j'}$, $x_{j''}$) belong to the same tuple τ_i (resp. τ_j , $i \neq j$), therefore, $x_{i'} = x_{i''}$ and $x_{j'} = x_{j''}$.

Now set $w = (i'i'')(j'j'')$ to be the product of two commuting transpositions. Then, $w \in W_u$ and $s_\beta = s_{w(\alpha)}$ and by Lemma 2.10 we obtain

Proposition 3.1. *For the type A, the closure of any double moment graph ${}^Q\mathcal{G}^P$ coincides with \mathcal{G}^P .*

Type D. We realize a root system of type D as a subset of vectors in \mathbb{R}^n

$$\Sigma = \{\pm(e_i + e_j), \pm(e_i - e_j) \mid i \neq j, i, j = 1, \dots, n\},$$

where e_1, \dots, e_n are the standard basis vectors. We fix a set of simple roots

$$\Theta = \{\alpha_1 = e_1 - e_2, \dots, \alpha_{n-1} = e_{n-1} - e_n, \alpha_n = e_{n-1} + e_n\}.$$

The Weyl group W acts by permutations and even sign changes, i.e., reflections are given either by usual transpositions or by signed transpositions

$$(ij): e_i \mapsto e_j, e_j \mapsto e_i \quad (\widetilde{ij}): e_i \mapsto -e_j, e_j \mapsto -e_i.$$

Consider the vector $\mu = u(\theta) \in W\theta$, where θ is dominant with stabilizer W_P and $u \in {}^QW^P$. Suppose $\widehat{s_\alpha u} = \widehat{s_\beta u} > u$ for some $\alpha, \beta \in \Sigma^+$.

Suppose $\alpha_n \notin \Theta_Q$. Then W_Q is a subgroup of the symmetric group S_n acting by permutations on $\{e_1, \dots, e_n\}$. Hence, as in type A , it acts on a vector $\mu = (x_1, \dots, x_n)$ by permutations of coordinates in each of the tuples (x_1, \dots, x_{k_1}) , $(x_{k_1+1}, \dots, x_{k_2})$, \dots , (x_{k_r+1}, \dots, x_n) , where $\Theta \setminus (\Theta_Q \cup \{\alpha_n\}) = \{\alpha_{k_1}, \dots, \alpha_{k_r}\}$. We have the following cases for the reflection s_α :

- (1) Let $s_\alpha = (i'j')$, $i' < j'$. As in type A , the coordinates $x_{i'}$ and $x_{j'}$ belong to different tuples and $x_{i'} \neq x_{j'}$.

- (2) Let $s_\alpha = \widetilde{(i'j')}$, $i' < j'$. Since $s_\alpha(\mu) \neq \mu$, we have $x_{i'} \neq -x_{j'}$ and
- (a) $x_{i'}$ and $x_{j'}$ belong to different tuples.
 - (b) $x_{i'}$ and $x_{j'}$ belong to the same tuple.

And we have similar cases (1), (2a) or (2b) for $s_\beta = (i''j'')$ or $\widetilde{(i'',j'')}$. Since $\widehat{s_\alpha u} = \widehat{s_\beta u}$ and the action by W_Q preserves the numbers of equal coordinates in each tuple, s_β corresponds to the same case as s_α and it interchanges coordinates (changes sign) between the same tuples as s_α , therefore, $x_{i'} = x_{i''}$ and $x_{j'} = x_{j''}$ (in the case (2b) we may permute i'' and j''). Taking $w = (i'i'')(j'j'')$ as before we obtain that $w \in W_u$ and $s_\beta = s_{w(\alpha)}$.

Suppose $\alpha_{n-1}, \alpha_n \in \Theta_Q$. The subgroup W_Q acts on the vector $\mu = (x_1, \dots, x_n)$ by permutations of coordinates in the first r tuples (x_1, \dots, x_{k_1}) , $(x_{k_1+1}, \dots, x_{k_2})$, \dots , $(x_{k_{r-1}+1}, \dots, x_{k_r})$, and by permutations and even sign changes on the last tuple $\tau = (x_{k_r+1}, \dots, x_n)$, where $\Theta \setminus \Theta_Q = \{\alpha_{k_1}, \dots, \alpha_{k_r}\}$. We have the following cases for the reflection s_α :

- (1) Let $s_\alpha = (i'j')$, $i' < j'$. As before, $x_{i'} \neq x_{j'}$ and $x_{i'}, x_{j'}$ belong to different tuples. We then have two subcases
 - (a) $x_{j'}$ does not belong to τ .
 - (b) $x_{j'}$ belongs to τ .
- (2) Let $s_\alpha = \widetilde{(i'j')}$, $i' < j'$. We have $x_{i'} \neq -x_{j'}$ and
 - (a) $x_{i'}$ and $x_{j'}$ belong to different tuples and $x_{j'}$ does not belong to τ .
 - (b) $x_{i'}$ and $x_{j'}$ belong to different tuples and $x_{j'}$ belongs to τ .
 - (c) $x_{i'}$ and $x_{j'}$ belong to the same tuple which is not τ .

And we have similar cases for $s_\beta = (i''j'')$ or $\widetilde{(i'',j'')}$. Observe that if both s_α and s_β correspond to the same case, then they interchange coordinates (signs) between the same tuples, hence, we can take the same w as before.

Since $\widehat{s_\alpha u} = \widehat{s_\beta u}$ and the action by W_Q preserves the numbers of equal coordinates in the first r tuples, s_α and s_β either both correspond to the case (1a), or they both correspond to the case (2a), or they both correspond to the remaining cases. Hence, we reduce to the situation when s_α, s_β correspond to different cases among (1b), (2b) or (2c).

Suppose s_α corresponds to (1b) and s_β corresponds to (2b), then we can take $w = (i'j')(i''j'')$ which is in W_u and $\beta = w(\alpha)$.

Suppose s_α corresponds either to (1b) or (2b) and s_β corresponds to (2c). Without loss of generality we may assume $x_{j''} = 0$, which implies $x_{i'} = x_{i''} = -x_{j'} \neq 0$ for (1b) and $x_{i'} = x_{i''} = x_{j'} \neq 0$ for (2b). If τ does not have zeros as coordinates, then $s_\alpha(\mu)$ and $s_\beta(\mu)$ belong to different W_Q -orbits, a contradiction. Therefore, τ must contain a zero coordinate.

Observe that s_α and s_β are not W_u -conjugate (if they are conjugate by an element of W_Q , then j', j'' have to be in the same tuple as well as i', i''). Moreover, if τ contains a zero coordinate and $x_{j''} = 0$, $x_{i'} = x_{i''} = -x_{j'} \neq 0$ (resp. $x_{j''} = 0$, $x_{i'} = x_{i''} = x_{j'} \neq 0$), then $(i'j')(\mu)$ (resp. $\widetilde{(i'j')}(\mu)$) and $(i'',j'')(\mu)$ belong to the same W_Q -orbit but are not W_Q -conjugate.

These arguments allow us to prove the following

Proposition 3.2. *For the type D , the closure of a double moment graph ${}^Q\mathcal{G}^P$ does not coincide with \mathcal{G}^P if and only if both $\Theta' = \Theta_P$ and $\Theta' = \Theta_Q$ satisfy*

- (i) $\alpha_{n-1}, \alpha_n \in \Theta'$ and (ii) $\Theta' \neq \{\alpha_s, \alpha_{s+1}, \dots, \alpha_n\}$ for any $s \leq n-1$.

Proof. If condition (i) for Θ_Q holds, then the tuple τ above contains at least two coordinates. Condition (ii) for Θ_Q says that some other tuple contains at least two coordinates, so the case (2c) above may appear. Condition (i) for Θ_P is equivalent to the condition that μ has at least two zero coordinates. Permuting coordinates we may assume that one zero coordinate belongs to the last tuple τ and another one belongs to a different tuple τ' . Condition (ii) for Θ_P means that μ contains two non-zero coordinates such that one belongs to τ and another one to τ' . \square

Type B . We realize a root system of type B as a subset of vectors in \mathbb{R}^n

$$\Sigma = \{\pm(e_i + e_j), \pm(e_i - e_j), \pm e_i \mid i \neq j, i, j = 1, \dots, n\},$$

where e_1, \dots, e_n are the standard basis vectors. We fix a set of simple roots

$$\Theta = \{\alpha_1 = e_1 - e_2, \dots, \alpha_{n-1} = e_{n-1} - e_n, \alpha_n = e_n\}.$$

The Weyl group W acts by permutations and sign changes, i.e., reflections are given by usual transpositions, by signed transpositions, and by sign changes

$$(ij): e_i \mapsto e_j, e_j \mapsto e_i \quad (\widetilde{ij}): e_i \mapsto -e_j, e_j \mapsto -e_i, \quad (\widetilde{i}): e_i \mapsto -e_i.$$

Example 3.3. For the Weyl group $W = \langle s_1 = (12), s_2 = (\widetilde{2}) \rangle$ of type B_2 set $\Theta_P = \{\alpha_2\}$ and $\Theta_Q = \{\alpha_1\}$. Then

$$W^P = \{e, s_1, s_2 s_1, s_1 s_2 s_1\}.$$

Set $\theta = e_1 = \alpha_1 + \alpha_2$ to be the dominant vector with stabilizer W_P . Then in terms of W -orbits we get

$$W^P = \{(1, 0), (0, 1), (0, -1), (-1, 0)\}.$$

As for double cosets we obtain ${}^QW^P = \{e, s_2 s_1\} = \{(1, 0), (0, -1)\}$. We have

$$\widehat{s_{e_1+e_2}} = \widehat{s_2 s_1 s_2} = s_2 s_1 \text{ and } \widehat{s_{e_1}} = \widehat{s_1 s_2 s_1} = s_2 s_1.$$

Observe that the roots e_1 and $e_1 + e_2$ belong to different W_Q -orbits.

Similar arguments as in type D imply the following type B version of Proposition 3.2.

Proposition 3.4. *For the type B the closure of a double moment graph ${}^Q\mathcal{G}^P$ coincides with \mathcal{G}^P if and only if one of the following is satisfied (we assume a type B_1 subsystem is generated by α_n only)*

- Both Θ_P and Θ_Q are subsystems of type B ,
- Both Θ_P and Θ_Q do not contain a type B subsystem,
- Either Θ_P or Θ_Q coincides with Θ .

4. EQUIVARIANT ALGEBRAIC ORIENTED COHOMOLOGY

Formal group laws. Let R be a commutative unital ring. Consider a one-dimensional commutative formal group law F over R that is a power series in two variables $F(x, y) \in R[[x, y]]$ which satisfies $F(x, y) = F(y, x)$, $F(x, 0) = x$, $F(F(x, y), z) = F(x, F(y, z))$. It can be written as

$$F(x, y) = x + y + \sum_{i, j > 0} a_{ij} x^i y^j, \text{ for some coefficients } a_{ij} \in R.$$

There is a formal inverse $-_F x$ that is a power series in one variable satisfying $F(x, -_F x) = 0$. The ring generated by all a_{ij} 's modulo relations imposed by commutativity and associativity is called the Lazard ring \mathbb{L} . There is a universal formal group law F_U over \mathbb{L} together with the evaluation map $F_U \rightarrow F$ (given by evaluating the coefficients). We set $x +_F y = F(x, y)$.

Example 4.1. Basic examples of formal group laws (f.g.l.) are

The additive f.g.l. $x +_F y = x + y$ over $R = \mathbb{Z}$ with $F_U \rightarrow F$ given by $a_{ij} \mapsto 0$ and $-_F x = -x$.

The multiplicative f.g.l. $x +_F y = x + y - \beta xy$ over $R = \mathbb{Z}[\beta]$ with $F_U \rightarrow F$ given by $a_{11} \mapsto -\beta$, $a_{ij} \mapsto 0$ for all $i, j > 1$ and $-_F x = \frac{x}{\beta x - 1}$. If β is invertible and $R = \mathbb{Z}[\beta^{\pm 1}]$, then the respective F is called multiplicative periodic.

Equivariant cohomology theories. We fix a base field k of characteristic 0. Let G be a split semisimple (not necessarily simply-connected) linear algebraic group over k . Consider a category of G -varieties over k with G -equivariant maps as morphisms. We denote by \mathbf{h}_G a G -equivariant algebraic oriented Borel-Moore homology theory introduced by Heller-Malagon-Lopez in [HM13] (it is obtained from a usual oriented Borel-Moore homology theory \mathbf{h} of Levine-Morel [LM07] generalizing the Totaro construction [To99]).

A key property of \mathbf{h} and, hence, of \mathbf{h}_G is that it has characteristic classes which satisfy the Quillen formula for the tensor product of line bundles [LM07, Lemma 1.1.3]. Namely,

$$c_1^{\mathbf{h}}(\mathcal{L}_1 \otimes \mathcal{L}_2) = F(c_1^{\mathbf{h}}(\mathcal{L}_1), c_1^{\mathbf{h}}(\mathcal{L}_2)),$$

where c_1 is the first characteristic class in the theory \mathbf{h} , \mathcal{L}_i is a line bundle over some variety X and $F(x, y) \in R[[x, y]]$ is the associated formal group law over $R = \mathbf{h}(pt)$ (here $pt = Spec k$).

There is a correspondence between \mathbf{h} 's and F 's: given \mathbf{h} one associates the respective F from the Quillen formula and given F over R one defines

$$\mathbf{h}(-) := \Omega(-) \otimes_{\Omega(pt)} R,$$

where Ω is the universal oriented theory (algebraic cobordism of Levine-Morel) over the $\Omega(pt)$. Observe that theories obtained in such a way satisfy short localization sequence

$$\mathbf{h}(Z) \rightarrow \mathbf{h}(X) \rightarrow \mathbf{h}(U) \rightarrow 0,$$

where U is open in X and $Z = X \setminus U$.

Example 4.2. The Chow theory $\mathbf{h} = CH$ corresponds to an additive formal group law $F(x, y) = x + y$ over $R = \mathbb{Z}$ [LM07, Ex.1.1.4] and CH_G is the G -equivariant Chow group in the sense of Totaro [To99, EG98].

Example 4.3. Following [LM07, Ex.1.1.4] consider a Grothendieck K^0 . Let

$$K^0(X)[\beta^{\pm 1}] = K^0(X) \otimes_{\mathbb{Z}} \mathbb{Z}[\beta^{\pm 1}]$$

be its \mathbb{Z} -graded version, where β is the Bott element (here we assign $\deg \beta = -1$ and $\deg x = 0$ for all $x \in K^0(X)$). Then $K^0(X)[\beta^{\pm 1}]$ is an oriented theory in the sense of [LM07] which corresponds to the multiplicative periodic formal group law $F(x, y) = x + y - \beta xy$ over $R = \mathbb{Z}[\beta^{\pm 1}]$ and $K_G^0(X)$ is the equivariant K -theory. If we set $\beta = 1$, then the \mathbb{Z} -graded version of K^0 is the direct sum of copies of $K^0(X)$ with each copy assigned the respective degree.

Example 4.4. If F is a formal group law of an elliptic curve, then it was shown in [ZZ17] that the associated oriented cohomology theory \mathfrak{h}_G is the local version of the equivariant elliptic cohomology constructed by Ginzburg-Kapranov-Vasserot [GKV].

Example 4.5. The G -equivariant algebraic cobordism Ω_G constructed in [HM13] corresponds to a universal formal group law F_U over the Lazard ring \mathbb{L} .

Equivariant coefficient rings. Suppose that $G = T$ is a split torus. Consider the character group T^* . Following [CPZ, §2] consider a completion $R[[x_\lambda]]_{\lambda \in T^*}$ of the polynomial ring $R[x_\lambda]_{\lambda \in T^*}$ in variables x_λ with respect to the kernel of the augmentation map $x_\lambda \mapsto 0$. Set

$$S = R[[x_\lambda]]_{\lambda \in T^*} / J$$

where J is the closure of the ideal of relations $x_0 = 0$ and $x_{\lambda+\mu} = x_\lambda +_F x_\mu$ for all $\lambda, \mu \in T^*$. The ring S is called a formal group algebra associated to G and F . According to [CZZ2, §3], the R -algebra S is the completion of the T -equivariant coefficient ring $\mathfrak{h}_T(pt)$ along the kernel of the forgetful map $\mathfrak{h}_T(pt) \rightarrow R$, i.e.

$$S \simeq \mathfrak{h}_T(pt)^\wedge.$$

Under this identification x_λ is the first characteristic class $c_1^{\mathfrak{h}}(\mathcal{L}_\lambda)$ of the associated line bundle \mathcal{L}_λ .

Example 4.6. For the additive f.g.l., i.e., for the Chow theory, if $G = PGL_2$, then $S = \mathbb{Z}[[\alpha]]$ where α is a simple root. If $G = SL_2$, then $S = \mathbb{Z}[[\omega]]$ where ω is a fundamental weight ($\alpha = 2\omega$).

Example 4.7. For the multiplicative periodic f.g.l., i.e. for $K^0(-)[\beta^{\pm 1}]$, the formal group algebra is the completed group ring $R[[T^*]]^\wedge$ (here we identify x_λ with $\beta^{-1}(1 - e^{-\lambda})$). For instance, if $G = SL_2$, then $S = R[[t^{\pm 1}]]^\wedge$, where $t = e^\omega$ and the completion is taken with respect to the kernel of $R[[t^{\pm 1}]] \rightarrow R$, $t \mapsto 1$.

Remark 4.8. If the formal group law F is a polynomial (e.g. for CH or K^0) and X is a projective homogeneous G -variety, in all subsequent construction we may substitute S by its non-completed version (the polynomial ring $S(T^*)$ and the group ring $\mathbb{Z}[[T^*]]$ respectively) as it does not affect the product structure in cohomology.

5. ORIENTED COHOMOLOGY SHEAVES ON MOMENT GRAPHS

Oriented sheaves. Let G be a split semisimple linear algebraic group over k , let T be its split maximal torus. Let $\mathfrak{h}_G(-)$ (resp. $\mathfrak{h}_T(-)$) be a G -equivariant (resp. T -equivariant) algebraic oriented cohomology theory. Let $\Sigma = \Sigma^+ \amalg \Sigma^- \supset \Theta$ be

the associated root datum and let $\mathcal{G} = ((V, \leq), l: E \rightarrow \Sigma^+)$ be a moment graph. For an edge $x \rightarrow y \in E$, we will write $x \text{ --- } y$ if we consider it only as an unordered pair of vertices. Observe that any root (character) $\alpha: T^* \rightarrow \mathbb{G}_m$ corresponds to a T -equivariant line bundle \mathcal{L}_α over pt and, hence, it gives the characteristic class $c_1^h(\mathcal{L}_\alpha) \in \mathfrak{h}_T(pt)$.

The following definition extends to any equivariant algebraic oriented cohomology theory the notion of moment graph sheaves from [BMP].

Definition 5.1. An \mathfrak{h} -sheaf \mathcal{F} on \mathcal{G} is given by the data $(\{\mathcal{F}^x\}, \{\mathcal{F}^{x-y}\}, \{\rho_{x,x-y}\})$, where:

- (SH1) for all $x \in V$, \mathcal{F}^x is an $\mathfrak{h}_G(pt)$ -module,
- (SH2) for all $x \text{ --- } y \in E$, \mathcal{F}^{x-y} is an $\mathfrak{h}_T(pt)$ -module such that

$$c_1^h(\mathcal{L}_{l(x-y)}) \cdot \mathcal{F}^{x-y} = (0),$$

i.e. it is annihilated by the respective T -equivariant characteristic class in the theory \mathfrak{h} ,

- (SH3) for $x \in V$ and $x \text{ --- } y \in E$, the map $\rho_{x,E}: \mathcal{F}^x \rightarrow \mathcal{F}^{x-y}$ is a homomorphism of $\mathfrak{h}_G(pt)$ -modules.

Example 5.2. Let $\mathfrak{h} = CH(-; \mathbb{Q})$ be the Chow theory with rational coefficients. Then $\mathfrak{h}_T(pt)$ is the polynomial ring in simple roots over \mathbb{Q} denoted simply by S . The Weyl group W of G acts naturally on S . Its subring of invariants S^W can be identified with $\mathfrak{h}_G(pt)$. The following data define an $CH(-; \mathbb{Q})$ -sheaf on the parabolic moment graph \mathcal{G}^P of Example 2.3:

- (SH1) $\mathcal{F}^x = S$, for all $x \in W^P$,
- (SH2) for all $x, y \in W^P$, $y = \overline{s_\alpha x}$, $\alpha \in \Sigma^+$, $\mathcal{F}^{x-y} = S/\alpha S$,
- (SH3) each map $\rho_{x,E}$ is simply the quotient $S \rightarrow S/\alpha S$.

This sheaf is called *sheaf of rings* in [BMP], and *structure sheaf* in [Fi08] and subsequent papers.

Example 5.3. Let $\mathfrak{h} = K^0(-)[\beta^{\pm 1}]$, $\beta = 1$ be the K -theory. Then $\mathfrak{h}_T(pt)$ is the group ring $Z = \mathbb{Z}[T^*]$ in characters of T . The subring of invariants Z^W can be identified with $\mathfrak{h}_G(pt)$. Each root α defines the characteristic class $c_1^h(\mathcal{L}_\alpha) = 1 - e^{-\alpha}$. The following data define an K^0 -sheaf on \mathcal{G}^P :

- (SH1) $\mathcal{F}^x = Z$, for all $x \in W^P$,
- (SH2) for all $x, y \in W^P$, $y = \overline{s_\alpha x}$, $\alpha \in \Sigma^+$, $\mathcal{F}^{x-y} = Z/(1 - e^{-\alpha})Z$,
- (SH3) each map $\rho_{x,E}$ is simply the quotient $Z \rightarrow Z/(1 - e^{-\alpha})Z$.

Oriented structure sheaves. More generally, we have

Example 5.4. For an arbitrary oriented cohomology theory \mathfrak{h} the following data define an \mathfrak{h} -sheaf \mathcal{F} on \mathcal{G}^P :

- (SH1) for all $x \in W^P$, we set $\mathcal{F}^x := S$, where $S = \mathfrak{h}_T(pt)^\wedge$;
- (SH2) for all $x \text{ --- } x' \in E$, we set $\mathcal{F}^{x-x'} := S/x_{l(x-x')}S$;
- (SH3) for $x \in W^P$ and $x \rightarrow x' \in E$, we set $\rho_{x,x-x'}$ to be the quotient map $S \rightarrow S/x_{l(x-x')}S$;

We call such \mathcal{F} a (parabolic) structure \mathfrak{h} -sheaf on \mathcal{G}^P and denote it by \mathcal{O}^P .

Extending to double moment graphs we obtain

Example 5.5. In the notation of Section 2 consider a double moment graph

$${}^Q\mathcal{G}^P = (({}^QW^P, \leq), l: {}^QE^P \rightarrow \Sigma^+)$$

for G , P and Q . The following data define an \mathfrak{h} -sheaf \mathcal{F} on ${}^Q\mathcal{G}^P$:

- (SH1) for all $u \in {}^QW^P$, we set $\mathcal{F}^u := S^{W_u}$, where $S = \mathfrak{h}_T(pt)^\wedge$;
- (SH2) for all $u \text{ --- } u' \in {}^QE^P$, we set $\mathcal{F}^{u \text{---} u'} := S/c_1^{\mathfrak{h}}(\mathcal{L}_{l(u \text{---} u')})S$;
- (SH3a) for $u \in {}^QW^P$ and $u \rightarrow u' \in {}^QE^P$, we set

$$\rho_{u, u \rightarrow u'} := \text{can} \circ \iota: \mathcal{F}^u \rightarrow \mathcal{F}^{u \rightarrow u'},$$

where $\iota: S^{W_u} \hookrightarrow S$ is the inclusion, and $\text{can}: S \rightarrow S/c_1^{\mathfrak{h}}(\mathcal{L}_{l(u \rightarrow u')})S$ is the canonical quotient map;

- (SH3b) Since $u' = s_{l(u \rightarrow u')}u \iff s_{l(u \rightarrow u')}u \in wu'W_P$, $w \in W_Q^{u'}$, we set

$$\rho_{u', u \rightarrow u'} := \text{can} \circ \iota_w: \mathcal{F}^{u'} \rightarrow \mathcal{F}^{u \rightarrow u'},$$

where $\iota_w: S^{W_{u'}} \hookrightarrow S$, is a twisted inclusion $s \mapsto w(s)$.

We call such \mathcal{F} a (double) structure \mathfrak{h} -sheaf and denote it ${}^Q\mathcal{O}^P$.

The following is a natural generalization to our setting of the module of local sections of a moment graph sheaf from [BMP, §1.3].

Definition 5.6. Let I be a subset of V , the module of local sections of an \mathfrak{h} -sheaf \mathcal{F} on a moment graph \mathcal{G} is defined as

$$\Gamma(I, \mathcal{F}) := \left\{ (m_x) \in \prod_{x \in I} \mathcal{F}^x \mid \begin{array}{l} \rho_{x, x-y}(m_x) = \rho_{y, y-x}(m_y) \\ \forall x \text{ --- } y \in E, x, y \in I \end{array} \right\}.$$

We write $\Gamma(\mathcal{F})$ for the module of global sections $\Gamma(V, \mathcal{F})$ of the \mathfrak{h} -sheaf \mathcal{F} .

Example 5.7. For the structure $CH(-; \mathbb{Q})$ -sheaf \mathcal{F} we have $\Gamma(\mathcal{F}) \simeq CH_T(G/P; \mathbb{Q})$ and for the structure K^0 -sheaf \mathcal{F} we have $\Gamma(\mathcal{F}) \simeq K_T^0(G/P)$. By [CZZ, Thm. 11.9] for an arbitrary \mathfrak{h} we have

$$\Gamma(\mathcal{O}^{\mathfrak{h}}) \simeq \mathfrak{h}_T(G/P).$$

Later we will show (see Theorem 9.1.(B)) that if $\langle {}^Q\mathcal{G}^P \rangle = \mathcal{G}^P$, then

$$\Gamma({}^Q\mathcal{O}^P) \simeq \mathfrak{h}_T(G/Q \times G/P)^W.$$

Equivariant oriented sheaves. We define an equivariant moment sheaf as follows

Definition 5.8. Given a W -equivariant moment graph \mathcal{G} of Definition 2.5, we say that an \mathfrak{h} -sheaf \mathcal{F} on \mathcal{G} is W -equivariant if it satisfies the following condition

- (SH4) for all $x \in V$, W acts by $\mathfrak{h}_G(pt)$ -linear automorphisms on \mathcal{F}^x and for each $w \in W$ there is an isomorphism of $\mathfrak{h}_G(pt)$ -modules $w(\mathcal{F}^{w^{-1}x}) \simeq \mathcal{F}^x$.

Definition 5.9. Let \mathcal{G} be a W -equivariant moment graph and let \mathcal{F} be a W -equivariant \mathfrak{h} -sheaf on \mathcal{G} . Let I be a W -invariant subset of V . The module of W -invariant local sections of \mathcal{F} is set to be

$$\Gamma(I, \mathcal{F})^W := \{(m_x) \in \Gamma(I, \mathcal{F}) \mid w(m_x) = m_{w(x)} \forall w \in W\}.$$

Remark 5.10. The study of the module of invariant local sections of the so-called BMP-sheaf (see, for example, [La12, §4]) on a Bruhat graph (equivariant with respect to an action of a Coxeter group) is a fundamental step in [La12, §6] towards the categorification of certain equalities of Kazhdan-Lusztig polynomials.

Example 5.11. Since W_Q acts on S via the action of the Weyl group on $\mathfrak{h}_T(pt)$, the parabolic structure sheaf \mathcal{O}^P is W_Q -equivariant. Later we will show (see Theorem 9.1.(A)) that

$$\Gamma(\mathcal{O}^P)^{W_Q} \simeq \mathfrak{h}_T(G/Q \times G/P)^W.$$

6. ORIENTED COHOMOLOGY OF FLAG VARIETIES

Let G be a split semi-simple linear algebraic group over k . We choose a split maximal torus T inside G . Let B be a Borel subgroup of G containing T . Let F be a formal group law over R corresponding to \mathfrak{h} .

The Borel case. Consider a complete flag variety G/B . We recall a construction of the T -equivariant oriented cohomology $\mathfrak{h}_T(G/B)$ which depends on the root system Σ of G , the character group T^* of T and the formal group law F of the theory \mathfrak{h} following [HMSZ, CZZ] (which is an extension of the arguments by Kostant-Kumar [KK86, KK90] to the context of an arbitrary \mathfrak{h}).

First, we define a twisted group algebra of the Weyl group W as the smash product $S_W = S \# R[W]$, where $S = \mathfrak{h}_T(pt)^\wedge$. By definition, S_W is a free left S -module with basis δ_w , $w \in W$ and multiplication satisfying $w(q)\delta_w = \delta_w q$. The S -linear dual S_W^* can be identified with the T -equivariant oriented cohomology of the T -fixed point set, i.e.,

$$S_W^* = \text{Hom}_S(S_W, S) = \text{Hom}(W, S) \simeq \mathfrak{h}_T((G/B)^T) = \bigoplus_{w \in W} \mathfrak{h}_T(pt_w).$$

Observe that S_W^* is a commutative ring where the product is given by the usual point-wise multiplication of functions $W \rightarrow S$ and at the same time it is induced by a cocommutative coproduct $\Delta: S_W \rightarrow S_W \otimes S_W$, $q\delta_w \mapsto q\delta_w \otimes \delta_w$.

Let $Q = S[\frac{1}{x_\alpha}, \alpha \in \Sigma]$ be the localization of S at all roots and let $Q_W = Q \# R[W]$ be the respective twisted group algebra. Following [HMSZ] we define the formal affine Demazure algebra \mathbb{D}_F to be the subalgebra of Q_W generated by push-pull elements

$$Y_i = \frac{1}{x_{-\alpha_i}} + \frac{1}{x_{\alpha_i}} \delta_{s_i},$$

for all simple roots α_i and multiplications by elements of S . The constant part of \mathbb{D}_F , i.e. the subalgebra generated only by Y_i 's over R , will be denoted by D_F .

Example 6.1. For the Chow theory, D_F (resp. \mathbb{D}_F) is a nil-Coxeter (resp. affine nil-Hecke) algebra. For the Grothendieck K^0 , D_F (resp. \mathbb{D}_F) is a (resp. affine) 0-Hecke algebra.

Finally, there is a ring isomorphism $\mathfrak{h}_T(G/B) \simeq \mathbb{D}_F^*$ between the T -equivariant oriented cohomology and the S -dual of \mathbb{D}_F . The natural inclusion $\eta: S_W \rightarrow \mathbb{D}_F$ induces the inclusion $\eta^*: \mathbb{D}_F^* \rightarrow S_W^*$ which is the restriction to the T -fixed point locus $\mathfrak{h}_T(G/B) \hookrightarrow \mathfrak{h}_T((G/B)^T)$.

The characteristic map. The algebra \mathbb{D}_F acts on its dual \mathbb{D}_F^* by means of the Hecke action ‘ \bullet ’, defined by (Q_W acts on Q_W^*)

$$q\delta_w \bullet pf_v = pvw^{-1}(q)f_{vw^{-1}}, \quad p, q \in Q,$$

where f_v is the standard basis (i.e., dual to δ_w) for Q -module Q_W^* . Under this action the elements Y_i of \mathbb{D}_F correspond to the classical push-pull operators

$$Y_i \bullet - : \mathfrak{h}_T(G/B) \rightarrow \mathfrak{h}_T(G/P_i) \rightarrow \mathfrak{h}_T(G/B),$$

where P_i is the minimal parabolic subgroup corresponding to the simple root α_i .

The equivariant characteristic map [CZZ1, Def. 6.8] $c: Q \rightarrow Q_W^*$ defined by $q \mapsto q \bullet \mathbf{1}$ restricts to $c: S \rightarrow \mathbb{D}_F^* = \mathfrak{h}_T(G/B)$. By definition, c is a W -equivariant R -algebra homomorphism given by $c(q) = \sum_{w \in W} w(q)f_w$. Consider the map

$$\rho: S \otimes_{S^W} S \rightarrow \mathfrak{h}_T(G/B), \quad (s, s') \mapsto s \cdot c(s').$$

The restriction to the T -fixed locus is an isomorphism if and only if the map ρ is an isomorphism (called the Borel isomorphism).

The Borel isomorphism holds if the ring of coefficients R contains \mathbb{Q} or if $\mathfrak{h} = K_0$ and G is simply-connected [St75]. For Chow groups with \mathbb{Z} -coefficients it fails in general (the image of η^* has a finite index).

The parabolic case. Let P be a parabolic subgroup of G and let G/P be the respective projective homogeneous variety. Let W_P denote the Weyl group of the Levi part of P . The above constructions can be extended to describe the T -equivariant oriented cohomology $\mathfrak{h}_T(G/P)$ as follows.

We first define $\mathbb{D}_{F,P}$ to be the image of \mathbb{D}_F under the composite $\mathbb{D}_F \rightarrow Q_W \rightarrow Q_{W/W_P}$, where Q_{W/W_P} is a free Q -module spanned by elements $\delta_{\bar{w}}$ corresponding to cosets $\bar{w} \in W/W_P$. Observe that $\mathbb{D}_{F,P}$ is not a ring but only a free left S -module. We set $\mathbb{D}_{F,P}^* = \text{Hom}_S(\mathbb{D}_{F,P}, S)$. The product on Q_W^* descends to the product on $\mathbb{D}_{F,P}^*$ and turns it into an algebra. The canonical map $\mathbb{D}_{F,P}^* \rightarrow \mathbb{D}_F^*$ induced by the projection $Q_W \rightarrow Q_{W/W_P}$ is injective, moreover, its image can be identified with the W_P -invariant subring via the Hecke action.

Finally, there are ring isomorphisms

$$\mathfrak{h}_T(G/P) \simeq (\mathbb{D}_F^*)^{W_P} \simeq \mathbb{D}_{F,P}^*.$$

The restriction to the T -fixed point locus $\mathfrak{h}_T(G/P) \rightarrow \mathfrak{h}_T((G/P)^T)$ is then induced by $S_{W/W_P} \rightarrow \mathbb{D}_{F,P}$. The Borel map restricts to W_P -invariants as well and turns into

$$\phi_P: S \otimes_{S^W} S^{W_P} \rightarrow \mathfrak{h}_T(G/P).$$

Therefore, if R contains \mathbb{Q} we can identify $\mathfrak{h}_T(G/P)$ with $S \otimes_{S^W} S^{W_P}$.

7. THE ORIENTED COHOMOLOGY OF THE PRODUCT

We now fix two parabolic subgroups P, Q of G , and denote by W_P , resp. W_Q , the Weyl groups of the Levi part of P , resp. Q . Our goal is to describe $\mathfrak{h}_G(G/Q \times G/P)$ as an $\mathfrak{h}_G(pt)$ -module generated by classes of orbits parametrized by double cosets.

We follow the notation of Section 2. For any $u \in QW^P$, we consider the parabolic subgroup P_u of G defined as $P_u := R_{unip}(Q) \cdot (Q \cap uPu^{-1})$. Then the Weyl group of its Levi part is $W_u = W_Q \cap uW_Pu^{-1}$.

Orbit stratification. Consider a diagonal action of G on the products of two projective homogeneous varieties $G/Q \times G/P$. Following [CM06, §2] the G -orbits in $G/Q \times G/P$ are parametrized by double cosets $W_Q \backslash W/W_P$. If $u \in {}^Q W^P$, then the orbit \mathcal{O}_u corresponding to u is the orbit of the pair (Q, uPu^{-1}) . The closure $\overline{\mathcal{O}}_w$ contains \mathcal{O}_v for another minimal double coset representative v if and only if $v \leq w$ in the Bruhat order on W .

For each orbit there is a G -equivariant affine fibration of rank $l(u)$ (see [CM06, §4])

$$\mathcal{O}_u \simeq G/(Q \cap uPu^{-1}) \rightarrow G/P_u.$$

In other words, $G/Q \times G/P$ has a G -equivariant relative cellular structure with a base consisting of varieties $\{G/P_u\}_{u \in {}^Q W^P}$.

By definition, the only closed orbit in $G/Q \times G/P$ corresponds to $u = e$ and if either P or Q is a Borel subgroup, then $P_u = B$ for any u .

Example 7.1. Consider a particular case $P = Q = B$. Then ${}^Q W^P = W$.

The closures of orbits $\overline{\mathcal{O}}_w$ admit G -equivariant resolutions of singularities $\widehat{\mathcal{O}}_w$ as follows (c.f. [NPSZ, §5]): Let $w = s_{i_1} s_{i_2} \dots s_{i_l}$ be a reduced expression as a product of simple reflections. Then the image of the projection to the first and the last factors

$$\widehat{\mathcal{O}}_w = G/B \times_{G/P_{i_1}} G/B \times_{G/P_{i_2}} \dots \times G/B \rightarrow G/B \times G/B$$

is the closure $\overline{\mathcal{O}}_w$.

Cohomology decomposition. The G -equivariant relative cellular structure on $G/B \times G/B$ then induces a direct sum decomposition of $\mathfrak{h}_G(pt)$ -modules

$$\mathfrak{h}_G(G/B \times G/B) \simeq \bigoplus_{w \in W} \mathfrak{h}_G(G/B)[\widehat{\mathcal{O}}_w] \simeq \bigoplus_{w \in W} S[\widehat{\mathcal{O}}_w].$$

For parabolic subgroups P and Q taking the projection of the same G -equivariant resolution $\widehat{\mathcal{O}}_u$ for a representative $u \in {}^Q W^P$ gives the resolution for the respective closure $\overline{\mathcal{O}}_u$, i.e.

$$\widehat{\mathcal{O}}_u \rightarrow G/B \times G/B \rightarrow G/Q \times G/P.$$

The G -equivariant cellular structure on $G/Q \times G/P$ then induces a direct sum decomposition of $\mathfrak{h}_G(pt)$ -modules

$$\mathfrak{h}_G(G/Q \times G/P) \simeq \bigoplus_{u \in {}^Q W^P} \mathfrak{h}_G(G/P_u)[\widehat{\mathcal{O}}_u],$$

i.e., any element x of $\mathfrak{h}_G(G/Q \times G/P)$ can be written uniquely as $x = \sum_{u \in {}^Q W^P} x_u$, where $x_u \in \mathfrak{h}_G(G/P_u)$. Observe that $\mathfrak{h}_G(G/Q \times G/P)$ is both $\mathfrak{h}_G(G/Q)$ -module and $\mathfrak{h}_G(G/P)$ -module via the pull-backs induced by projections $G/Q \times G/P \rightarrow G/Q, G/P$.

Consider the forgetful map $\mathfrak{h}_G(G/Q \times G/P) \rightarrow \mathfrak{h}_T(G/Q \times G/P)$. It is induced by restricting the coefficients $\mathfrak{h}_G(G/P_u) \rightarrow \mathfrak{h}_T(G/P_u)$ which is not necessarily injective. However, in some cases, e.g. if either $P = B$ or $Q = B \implies P_u = B$, we obtain an embedding $S \simeq \mathfrak{h}_G(G/B) \hookrightarrow \mathfrak{h}_T(G/B) \simeq \mathbb{D}_F^*$.

Let $\mathfrak{h}_{G|T}(X)$ denote the image of the forgetful map $\mathfrak{h}_G(X) \rightarrow \mathfrak{h}_T(X)$. Then $\mathfrak{h}_G(G/Q \times G/B) = \mathfrak{h}_{G|T}(G/Q \times G/B)$, $\mathfrak{h}_G(G/B \times G/P) = \mathfrak{h}_{G|T}(G/B \times G/P)$ and

$$\mathbf{h}_{G|T}(G/Q \times G/P) \simeq \bigoplus_{u \in {}^Q W^P} \mathbf{h}_{G|T}(G/P_u)[\widehat{\mathcal{O}}_u]$$

Remark 7.2. The case when $P = B$ or $Q = B$ is exactly the case when the base of the cellular filtration consists of one variety only that is G/B , i.e., $G/Q \times G/B$ or $G/B \times G/P$ is a G -equivariant cellular space over G/B via the second or the first projection. The latter implies that the respective cohomology is a free S -module.

Remark 7.3. There are examples of G , $P \neq B$ and $Q \neq B$ such that all maps $\mathbf{h}_G(G/P_u) \rightarrow \mathbf{h}_T(G/P_u)$ are injective, and $\mathbf{h}_G(G/Q \times G/P) = \mathbf{h}_{G|T}(G/Q \times G/P)$.

Since both G/Q and G/P are T -cellular spaces, by [NPSZ, §3] there is a Künneth isomorphism

$$(1) \quad \mathbf{h}_T(G/Q \times G/P) \simeq \mathrm{Hom}_{\mathbf{h}_T(pt)}(\mathbf{h}_T(G/Q), \mathbf{h}_T(G/P)),$$

given by $[\widehat{\mathcal{O}}_u] \mapsto Y_{I_u} \bullet -$, where $I_u = (i_1, \dots, i_l)$ is a reduced word for u and Y_{I_u} is the composite $Y_{i_1} \dots Y_{i_l}$ of push-pull operators.

8. THE W -ACTION ON THE PRODUCT

In the present section we extend arguments of [CNZ, §3] (where the case of $X = G/P$ was treated) to a product of two projective homogeneous varieties. We show that $\mathbf{h}_{G|T}(G/Q \times G/P)$ embeds into the W -invariants of $\mathbf{h}_T(G/Q \times G/P)$ in a way compatible with the orbit stratification (see Corollary 8.3).

Geometric W -action. Let X be an arbitrary (left) G -variety. Recall that there is a natural (left) action of W on $\mathbf{h}_T(X)$. It can be either realized by pull-backs induced by a right action of W on each step of the Borel construction

$$V \times^T X = V \times X / (v, x) \sim (vt, t^{-1}x), \quad t \in T$$

given by

$$(v, x)T \cdot \sigma T = (v\sigma, \sigma^{-1}x)T, \quad \sigma \in N_G(T)$$

where V is taken to have a right G -action; or through the natural isomorphism $\mathbf{h}_T(X) \simeq \mathbf{h}_G(G/T \times X)$ and a G -equivariant right action of W on the variety $G/T \times_k X$ given by the formula $(gT, x) \cdot \sigma T = (gT\sigma, x)$.

Now let $X = G/Q \times G/P$ be a product of projective homogeneous varieties where G acts diagonally. Consider the stratification of X into G -orbits \mathcal{O}_u , $u \in {}^Q W^P$. We identify the set of T -fixed points of the orbit \mathcal{O}_u with a finite constant scheme W/W_u with trivial T -action. In the induced pullback (the T -fixed point locus restriction)

$$\eta_{PQ}: \mathbf{h}_T(G/Q \times G/P) \rightarrow \bigoplus_{u \in {}^Q W^P} \mathbf{h}_T(W/W_u)$$

we identify $\mathbf{h}_T(W/W_u)$ with the ring of all set-theoretic maps $\mathrm{Maps}(W/W_u, \mathbf{h}_T(pt))$ via

$$\mathbf{h}(V \times^T W/W_u) = \bigoplus_{xW_u \in W/W_u} \mathbf{h}_T(pt) = \mathrm{Maps}(W/W_u, \mathbf{h}_T(pt)),$$

where the class of $(v, xW_u)T$ maps to $xW_u \mapsto [vT]$.

Lemma 8.1. *Consider the left W -action on $\text{Maps}(W/W_u, \mathfrak{h}_T(pt))$ given by*

$$(w \cdot f)(x) = w \cdot f(w^{-1}x), \quad x \in W/W_u, \quad f \in \text{Maps}(W/W_u, \mathfrak{h}_T(pt)).$$

Then the pullback map

$$\eta_{PQ}: \mathfrak{h}_T(G/Q \times G/P) \rightarrow \bigoplus_{u \in {}^Q W^P} \mathfrak{h}_T(W/W_u) = \bigoplus_{u \in {}^Q W^P} \text{Maps}(W/W_u, \mathfrak{h}_T(pt))$$

is W -equivariant.

Proof. Recall that the action of W on $U \times^T (G/Q \times G/P)$ is given by

$$(v, (gQ, gP))T \cdot \sigma T = (v\sigma, \sigma^{-1}(gQ, gP))T$$

for any V in the Borel construction. Restricted to $V \times^T W/W_u$ the action of $w = \sigma T$ gives a map

$$V \times^T W/W_u \rightarrow V \times^T W/W_u, \quad (v, xW_u)T = (v\sigma, \sigma^{-1}xW_u)T.$$

So its pullback defines an endomorphism of $\mathfrak{h}_T(W/W_u) = \text{Maps}(W/W_u, \mathfrak{h}_T(pt))$ given by $f \mapsto w \cdot f$, where $w \cdot f: x \mapsto wf(w^{-1}x)$. \square

Lemma 8.2. *The map η_{PQ} factors through the orbit decomposition and is injective, i.e.,*

$$\eta_{PQ}: \mathfrak{h}_T(G/Q \times G/P) \simeq \bigoplus_{u \in {}^Q W^P} \mathfrak{h}_T(G/P_u)[\widehat{\mathcal{O}}_u] \xrightarrow{\oplus_u \eta_u} \bigoplus_{u \in {}^Q W^P} \mathfrak{h}_T(W/W_u)$$

where $\eta_u: \mathfrak{h}_T(G/P_u) \rightarrow \mathfrak{h}_T(W/W_u)$ is the T -fixed point locus restriction.

Proof. It is compatible with the orbit stratification by definition and it is injective since so is each η_u . \square

By definition of the W -action the forgetful map $\mathfrak{h}_G(pt) \rightarrow \mathfrak{h}_T(pt)$ factors through ${}^W \mathfrak{h}_T(pt)$ but the map $\mathfrak{h}_G(pt) \rightarrow {}^W \mathfrak{h}_T(pt)$ is neither injective nor surjective in general. Moreover, $\mathfrak{h}_{G|T}(pt)$ is of finite index in ${}^W \mathfrak{h}_T(pt) = S^W$.

Corollary 8.3. *There is an isomorphism of ${}^{W^P} \mathfrak{h}_T(G/P)$ - and ${}^{W^Q} \mathfrak{h}_T(G/Q)$ -modules*

$${}^W \mathfrak{h}_T(G/Q \times G/P) \simeq \bigoplus_{u \in {}^Q W^P} {}^{W^u} \mathfrak{h}_T(G/P_u)[\widehat{\mathcal{O}}_u]$$

and $\mathfrak{h}_{G|T}(G/Q \times G/P)$ is a subgroup of finite index in ${}^W \mathfrak{h}_T(G/Q \times G/P)$.

Proof. It is obviously an isomorphism of ${}^W \mathfrak{h}_T(G/B) = S^W$ -modules. The structure of ${}^{W^P} \mathfrak{h}_T(G/P)$ -module (resp. ${}^{W^Q} \mathfrak{h}_T(G/Q)$ -module) on ${}^{W^u} \mathfrak{h}_T(G/P_u)$ is induced by projection $G/P_u \rightarrow G/P$ (resp. $G/P_u \rightarrow G/Q$) and the isomorphism is compatible with these projections. \square

Example 8.4. In particular, if the Borel isomorphism holds for \mathfrak{h} (e.g. if R contains \mathbb{Q}), then $\mathfrak{h}_G(pt) = \mathfrak{h}_{G|T}(pt) = {}^W \mathfrak{h}_T(pt)$, ${}^{W^P} \mathfrak{h}_T(G/P) = S^{W^P}$, ${}^{W^Q} \mathfrak{h}_T(G/Q) = S^{W^Q}$ so

$$\mathfrak{h}_G(G/Q \times G/P) = \mathfrak{h}_{G|T}(G/Q \times G/P) = {}^W \mathfrak{h}_T(G/Q \times G/P)$$

and

$$\mathfrak{h}_G(G/Q \times G/P) \simeq S^{W^Q} \otimes_{S^W} S^{W^P}.$$

The map η_{PQ} then is an isomorphism

$$\eta_{PQ}: S^{W_Q} \otimes_{S^W} S^{W_P} \xrightarrow{\cong} \bigoplus_{u \in QW_P} S^{W_u}.$$

Algebraic W -action. Following [LZZ, §3] we introduce another Q_W action on Q_W^* by

$$(p\delta_w) \odot (qf_v) = pw(q)f_{wv}, \quad p, q \in Q, w, v \in W.$$

This action restricts exactly to an action of \mathbb{D}_F on $\mathbb{D}_F^* = \mathfrak{h}_T(G/B)$. Moreover, the respective action by $\delta_w \in \mathbb{D}_F$ gives the action by $w \in W$ on the cohomology $\mathfrak{h}_T(G/B)$, i.e., the W -action on $\mathfrak{h}_T(G/B)$ is given by the \odot -action.

Remark 8.5. The \odot -action appeared in the context of Chow groups and K -theory in the works by Tymoszko, Brion, Peterson, Knutson and others.

So we have two different actions on \mathbb{D}_F^* : the \odot -action (the left action) and the Hecke \bullet -action (the right action). There are two commutative diagrams

$$\begin{array}{ccc} S \otimes_{S^W} S & \xrightarrow{\rho} & \mathbb{D}_F^* \\ (z \cdot) \otimes 1 \downarrow & & \downarrow z \odot - \\ S \otimes_{S^W} S & \xrightarrow{\rho} & \mathbb{D}_F^* \end{array} \quad \begin{array}{ccc} S \otimes_{S^W} S & \xrightarrow{\rho} & \mathbb{D}_F^* \\ 1 \otimes (z \cdot) \downarrow & & \downarrow z \bullet - \\ S \otimes_{S^W} S & \xrightarrow{\rho} & \mathbb{D}_F^* \end{array}$$

where ρ is the Borel map and $z \in S_W$ acts on S by means of the \cdot -action. In other words, the \odot corresponds to the action of W on the left S -factor and \bullet corresponds to the action of W on the right S -factor in $S \otimes_{S^W} S$.

We know that $\mathfrak{h}_T(G/B)^{W_P} = (\mathbb{D}_F^*)^{W_P} \simeq \mathfrak{h}_T(G/P)$ (the \bullet -action of W_P). The respective Borel model of this isomorphism is then $(S \otimes_{S^W} S)^{W_P} \simeq S \otimes_{S^W} S^{W_P}$. We have ${}^{W_Q}(S \otimes_{S^W} S)^{W_P} \simeq S^{W_Q} \otimes_{S^W} S^{W_P}$ which corresponds to

$${}^{W_Q} \mathfrak{h}_T(G/B)^{W_P} \simeq {}^W \mathfrak{h}_T(G/Q \times G/P).$$

Since the Künneth isomorphism (1) is W -equivariant we obtain

$$\begin{aligned} {}^W \mathfrak{h}_T(G/Q \times G/P) &\simeq \text{Hom}_{W\text{-equiv}}(\mathfrak{h}_T(G/Q), \mathfrak{h}_T(G/P)) \\ &\simeq \text{Hom}_{S_W}((\mathbb{D}_F^*)^{W_Q}, (\mathbb{D}_F^*)^{W_P}) \\ &\simeq \text{Hom}_{S_W}(\mathbb{D}_{F,Q}^*, \mathbb{D}_{F,P}^*), \end{aligned}$$

where \mathbb{D}_F^* is considered as an S_W -module via the \odot -action.

9. GLOBAL SECTIONS OF STRUCTURE SHEAVES

In the present section we describe W -invariants of the T -equivariant cohomology of a product ${}^W \mathfrak{h}_T(G/Q \times G/P)$ as modules of global sections of structure sheaves on moment graphs. Our main result is the following

Theorem 9.1. *As an S^{W_Q} -module, ${}^W \mathfrak{h}_T(G/Q \times G/P)$ is isomorphic to*

$$(A) \quad {}^{W_Q} R_{W_P} = \left\{ (b_v) \in \bigoplus_{v \in W_P} S^{W_v} \left| \begin{array}{l} \text{(i) } b_v - b_{\overline{s_\beta v}} \in x_\beta S \text{ for all } s_\beta \in W, \text{ and} \\ \text{(ii) } b_{\overline{s_\alpha v}} = s_\alpha(b_v) \text{ for all } s_\alpha \in W_Q, \end{array} \right. \right\}$$

and, hence, it is isomorphic to the module of W_Q -invariant global sections $\Gamma(\mathcal{O}^P)^{W_Q}$ of the structure \mathfrak{h} -sheaf \mathcal{O}^P of Example 5.4.

$$(B) \quad {}^Q A^P := \left\{ (c_u) \in \bigoplus_{u \in {}^Q W^P} S^{W_{P_u}} \mid \begin{array}{l} c_u - w(c_{u'}) \in x_\alpha S \\ \text{for all } \alpha \in \Sigma^+, w \in W_Q \\ \text{s. t. } u \in s_\alpha w u' W_P \end{array} \right\}.$$

Moreover, if $\langle {}^Q \mathcal{G}^P \rangle = \mathcal{G}^P$, then it is isomorphic to the module of global sections $\Gamma({}^Q \mathcal{O}^P)$ of the double structure sheaf of Example 5.5.

Proof. (A) Observe that the map η_{PQ} restricted to W -invariants gives an inclusion (the T -fixed locus restriction)

$$\eta_{PQ}: {}^W \mathfrak{h}_T(G/Q \times G/P) \simeq \bigoplus_{u \in {}^Q W^P} W_u \mathfrak{h}_T(G/P_u) \xrightarrow{\oplus \eta_u} \bigoplus_{u \in {}^Q W^P} S^{W_u} \hookrightarrow \bigoplus_{u \in {}^Q W^P} Q^{W_u}.$$

Hence, we can identify ${}^W \mathfrak{h}_T(G/Q \times G/P) = {}^W Q \mathfrak{h}_T(G/P)$ with an S^{W_Q} -submodule of $\bigoplus_{u \in {}^Q W^P} S^{W_u}$.

Using localization [CZZ, Thm. 11.9], we can identify $\mathfrak{h}_T(G/P)$ with the following S -submodule of S_{W/W_P}^* :

$$(*) \quad R_{W_P} = \left\{ (b_v) \in \bigoplus_{v \in W^P} S \mid b_v - b_{\overline{s_\alpha v}} \in x_\beta S, \forall s_\beta \notin v W_P v^{-1} \right\}$$

Observe that the divisibility does not depend on a choice of representative of the coset $s_\alpha v$ as there are no multiple edges in the respective (parabolic) moment graph.

The group W acts on $Q_{W_P}^*$ via the \odot -action given by

$$w \odot b_v f_v = w(b_v) f_{wv} \text{ or, equivalently, } w \odot (b_v) = (w(b_{w^{-1}v})).$$

This induces an action of W on S_{W/W_P}^* via $w \odot b_v f_v = w(b_v) f_{\overline{wv}}$ which restricts to an action on $\mathfrak{h}_T(G/P) \subset S_{W/W_P}^*$.

Now, $(b_v) \in R_{W_P}$ is W_Q -invariant if and only if for any $v \in W^P$ it holds

$$(**) \quad b_{\overline{s_\alpha v}} = s_\alpha(b_v), \quad \forall s_\alpha \in W_Q.$$

Observe that $\overline{s_\alpha v} = v$ if $s_\alpha \in v W_P v^{-1}$. Hence, $(b_v) \in R_{W_P}$ is W_v -invariant if and only if

$$b_v = b_{\overline{s_\alpha v}} = s_\alpha(b_v), \quad \forall s_\alpha \in W_v, \text{ i.e., } b_v \in S^{W_{P_v}}.$$

Combining (*) and (**) we prove the isomorphism (A). The identification with the W_Q -invariant sections follows from the definition of $\Gamma(\mathcal{O}^P)^{W_Q}$.

(B) Observe that α is uniquely determined by the class of w modulo $W_{u'}$, i.e., if $u \in s_\alpha w u' W_P$ and $u \in s_\beta w' u' W_P$, where w and w' belong to the same coset in $W_Q/W_{u'}$, then $\alpha = \beta$.

We define a map

$$\psi: {}^Q A^P \rightarrow \bigoplus_{v \in W^P} S^{W_{P_v}}, \quad \psi(c_u) := (w(c_u))_{wu}, \quad w \in W_Q^u.$$

This map is clearly an injective S^{W_Q} -module homomorphism, so if we show that its image is ${}^W Q R_{W_P}$ we are done.

Let $v = wu \in W^P$ be a parabolic decomposition. Since $s_\alpha v = s_\alpha w u$, $s_\alpha \in W_Q$ is in the same double coset as v , $\overline{s_\alpha v} = w' u$ is the parabolic decomposition of $\overline{s_\alpha v}$, where w' is the minimal representative of the coset of $s_\alpha w$ in W_Q/W_u . We

then have $b_{\overline{s_\alpha v}} = w'(c_u) = s_\alpha(w(c_u)) = s_\alpha(b_v)$. In other words, the image of ψ is W_Q -invariant, i.e. (ii) holds.

Consider parabolic decompositions $v = wu$ and $\overline{s_\beta v} = w'u'$, where $u, u' \in {}^QW^P$, w is a minimal representative in W_Q/W_u and w' is a minimal representative in $W_Q/W_{u'}$. Then $b_v = w(c_u)$, $b_{\overline{s_\beta v}} = w'(c_{u'})$ and the condition (i) becomes equivalent to

$$w(c_u) - w'(c_{u'}) \in x_\beta S \text{ and, hence, to}$$

$$c_u - w^{-1}w'(c_{u'}) \in x_{w^{-1}(\beta)} S.$$

Since $w'u'z = s_\beta v = s_\beta wu$ for some $z \in W_P$, we obtain $u = w^{-1}s_\beta w'u'z = s_{w^{-1}(\beta)}w^{-1}w'u'z$. Substituting $w^{-1}w'$ by w we obtain the condition on (c_u) in the definition of ${}^QA^P$, Hence, (i) holds.

We have shown that $\text{Im}(\psi) \subseteq {}^{W_Q}R_{W^P}$.

Vice versa, given an element $(b_v)_{v \in W^P}$, one can obtain an element in ${}^QA^P$ by projecting onto the ${}^QW^P$ -components: Set $c_u = w^{-1}(b_v)$, where $v = wu$ is the parabolic decomposition. If $u, u' \in {}^QW^P$, $w \in W_Q$, and $\alpha \in \Sigma^+$ are such that $u \in s_\alpha wu'W_P$, then $c_u - w(c_{u'}) = c_u - c_{wu'} \in x_\alpha S$ (since $c_{u'} \in S^{W_{u'}}$ we may assume $wu' \in {}^QW^P$) and so $(c_u)_{u \in {}^QW^P} \in {}^QA^P$.

Finally, the identification with global sections $\Gamma({}^QO^P)$ follows by definition and the proof of part (B) is finished. \square

Example 9.2. Let $G = SL_3$ and let P_1 and P_2 denote the two parabolic subgroups

$$P_1 = \begin{pmatrix} * & * & * \\ * & * & * \\ 0 & 0 & * \end{pmatrix}, \quad P_2 = \begin{pmatrix} * & * & * \\ 0 & * & * \\ 0 & * & * \end{pmatrix}$$

Let $s_1 = (12)$, $s_2 = (23)$ be the two simple reflections of $W = S_3$ corresponding to the two parabolic subgroups above. Let $\mathfrak{h} = CH$ be the Chow theory. We then have $S := \mathbb{Z}[\alpha_1, \alpha_2]$ (the polynomial ring in two variables with integer coefficients).

The minimal length representatives for $S_3/\langle s_1 \rangle$ are $\{e, s_2, s_1s_2\}$ and the moment graph description of $CH_T(SL_3/P_1)$ is

$$\left\{ (z_e, z_{s_2}, z_{s_1s_2}) \in S \oplus S \oplus S \mid \begin{array}{l} z_e - z_{s_2} \in \alpha_2 S, \quad z_{s_2} - z_{s_1s_2} \in \alpha_1 S \\ z_e - z_{s_1s_2} \in (\alpha_1 + \alpha_2) S \end{array} \right\}.$$

Here $s_1s_2 = s_{s_2(\alpha_1 + \alpha_2)}s_2$, where $s_2(\alpha_1 + \alpha_2) = \alpha_1$.

The action of S_3 on $CH_T(SL_3/P_1)$ is determined by

$$(2) \quad s_1((z_e, z_{s_2}, z_{s_1s_2})) = (s_1(z_e), s_1(z_{s_1s_2}), s_1(z_{s_2})),$$

$$(3) \quad s_2((z_e, z_{s_2}, z_{s_1s_2})) = (s_2(z_{s_2}), s_2(z_e), s_2(z_{s_1s_2})).$$

By equation (2), an element $(z_e, z_{s_2}, z_{s_1s_2}) \in CH_T(SL_3/P_1)$ is s_1 -invariant if and only if

$$z_e = s_1(z_e), \quad z_{s_2} = s_1(z_{s_1s_2}).$$

(The third condition $z_{s_1s_2} = s_1(z_{s_2})$ is clearly redundant). We deduce that

$${}^{s_1}CH_T(SL_3/P_1) =$$

$$\{(z_e, z_{s_2}, s_1(z_{s_2})) \in S^{s_1} \oplus S \oplus S \mid z_e - z_{s_2} \in \alpha_2 S, z_e - s_1(z_{s_2}) \in (\alpha_1 + \alpha_2) S\}.$$

We notice that the second congruence is equivalent to the first one, since $z_e \in S^{s_1}$ and $s_1(\alpha_2) = \alpha_1 + \alpha_2$. Thus we have an isomorphism of S^{s_1} -modules:

$${}^{s_1}CH_T(SL_3/P_1) \cong \{(z_e, z_{s_2}) \in S^{s_1} \oplus S \mid z_e - z_{s_2} \in \alpha_2 S\}.$$

This description is encoded in the double moment graph:

$$e \xrightarrow{\alpha_2} s_2.$$

We notice that $\{(1, 1), (\alpha_2(\alpha_1 + \alpha_2), 0), (0, \alpha_2)\}$ is a basis for ${}^{s_1}CH_T(SL_3/P_1)$ as a free S^{s_1} -module.

Example 9.3. Equation (3) tells us that $(z_e, z_{s_2}, z_{s_1 s_2}) \in CH_T(SL_3/P_1)$ is s_2 -invariant if and only if

$$z_{s_2} = s_2(z_e) \quad \text{and} \quad z_{s_1 s_2} = s_2(z_{s_1 s_2}),$$

from which, reasoning as in the previous case, we get

$${}^{s_2}CH_T(SL_3/P_1) \cong \{(z_e, z_{s_1 s_2}) \in S \oplus S^{s_2} \mid z_e - z_{s_2 s_1} \in (\alpha_1 + \alpha_2)S\}.$$

Such a space is an S^{s_2} -module with a basis $\{(1, 1), (\alpha_1 + \alpha_2, 0), (0, \alpha_1(\alpha_1 + \alpha_2))\}$. The corresponding double moment graph is then

$$e \xrightarrow{\alpha_1 + \alpha_2} s_1 s_2.$$

10. CORRESPONDENCE PRODUCT AND EQUIVARIANT MOTIVES

Correspondence product. The $\mathfrak{h}_G(pt)$ -module $\mathfrak{h}_G(G/B \times G/B)$ can be endowed with another product structure the so called correspondence product. Namely, given $f, g \in \mathfrak{h}_G(G/B \times G/B)$ we set

$$f \circ g := pr_{1,3*}(pr_{1,2}^*(f) \cdot pr_{2,3}^*(g)),$$

where $pr_{i,j}: (G/B)^3 \rightarrow (G/B)^2$ are projections to the respective components and pr^* , pr_* denote the induced pull-backs and push-forwards in the theory \mathfrak{h} . Since $G/B \times G/B$ is a cellular space over G/B (via the orbit stratification), and $\mathfrak{h}_G(G/B) = \mathfrak{h}_T(pt) = {}^W\mathfrak{h}_T(G/B)$, we have

$$\mathfrak{h}_G(G/B \times G/B) = \mathfrak{h}_{G|T}(G/B \times G/B) = {}^W\mathfrak{h}_T(G/B \times G/B) = {}^B A^B$$

as $\mathfrak{h}_G(pt)$ -modules. By [NPSZ, Theorem 6.2] ${}^B A^B$ with respect to the correspondence product \circ is a (non-commutative) ring isomorphic to \mathbb{D}_F . Moreover, by the Künneth isomorphism it is also isomorphic to the endomorphism ring $End_{S_W}(\mathbb{D}_F^*)$ of S_W -modules (w.r. to the \odot -action).

More generally, given parabolic subgroups P, Q, H of G there is a $\mathfrak{h}_G(pt)$ -bilinear map

$$\circ: \mathfrak{h}_G(G/Q \times G/P) \otimes \mathfrak{h}_G(G/P \times G/H) \rightarrow \mathfrak{h}_G(G/Q \times G/H),$$

defined by the correspondence product

$$(\phi, \psi) \mapsto \psi \circ \phi = pr_{Q,H*}(pr_{P,H}^*(\psi) \cdot pr_{Q,P}^*(\phi)),$$

where $pr_{Q,P}: G/Q \times G/P \times G/H \rightarrow G/Q \times G/P$ is the corresponding projection. It restricts to the cohomology of the image $\mathfrak{h}_{G|T}(G/Q \times G/P)$ and, moreover, since the Künneth isomorphism is W -equivariant it induces a pairing

$$\begin{aligned} {}^Q A^P \otimes {}^P A^H &\rightarrow {}^Q A^H, \text{ where} \\ {}^Q A^P &= ({}^W\mathfrak{h}_T(G/Q \times G/P), \circ) \simeq End_{S_W}(\mathbb{D}_{F,Q}^*, \mathbb{D}_{F,P}^*). \end{aligned}$$

Equivariant motives. Let \mathfrak{M}_G denote a full additive subcategory of the category of G -equivariant \mathfrak{h} -motives generated by the motives of varieties G/P for all parabolic P (we refer to [PS16] for definitions and basic properties of this category). Recall only that morphisms in this category are cohomology groups $\mathfrak{h}_G(X \times Y)$, where G acts diagonally on the product of smooth projective G -varieties X and Y and the composition is given by the correspondence product. Following [CNZ, NPSZ] let $\mathfrak{M}_{G|T}$ denote a full additive subcategory of the respective category of $G|T$ -equivariant \mathfrak{h} -motives where the morphisms are given by images $\mathfrak{h}_{G|T}(X \times Y)$ of forgetful maps.

Definition 10.1. Given a category of projective homogeneous G -varieties we define a category of W -invariant T -equivariant motives ${}^W\mathfrak{M}_T$ as follows. We set morphisms to be elements of ${}^W\mathfrak{h}_T(X \times Y)$ and the composition to be the restriction of the correspondence product on \mathfrak{M}_T (observe that the correspondence product on \mathfrak{M}_T is W -equivariant).

By Theorem 9.1 the category ${}^W\mathfrak{M}_T$ is equivalent to the full additive subcategory of S_W -modules generated by $\mathbb{D}_{F,P}^*$ for all parabolics P . There are induced functors

$$\mathfrak{M}_G \xrightarrow{\mathcal{F}_1} \mathfrak{M}_{G|T} \xrightarrow{\mathcal{F}_2} {}^W\mathfrak{M}_T$$

defined on morphisms by

$$\mathfrak{h}_G(G/Q \times G/P) \rightarrow \mathfrak{h}_{G|T}(G/Q \times G/P) \hookrightarrow {}^W\mathfrak{h}_T(G/Q \times G/P) = {}^Q A^P$$

so the functor \mathcal{F}_2 is faithful by definition.

Remark 10.2. All these functors become equivalences if R contains \mathbb{Q} or if the torsion index of G is 1 or if G is simply-connected and $\mathfrak{h} = K_0$.

Remark 10.3. For some theories \mathfrak{h} (e.g. for CH and K_0) induced maps on endomorphisms rings have nilpotent kernels (this is essentially the Rost nilpotence theorem for \mathfrak{h} discussed in [CM06, CNZ]). So there is a lifting of idempotents with respect to \mathcal{F}_1 . In other words, if $\text{End}_{S_W}(\mathbb{D}_{F,P}^*) = {}^P A^P$ has no idempotents, then the G -equivariant \mathfrak{h} -motive of G/P is indecomposable.

Remark 10.4. For the Chow theory $CH(-; \mathbb{Z}/p\mathbb{Z})$, p is a prime, and some groups G (e.g. see [Vi07] for PGL_p), the functor \mathcal{F}_2 is an equivalence. The latter is closely related to surjectivity of the map $CH(BG) \rightarrow CH(BT)^W$. Indeed, if these maps are surjective for Levi-parts of all parabolic subgroups of G , including G itself, then \mathcal{F}_2 is an equivalence.

Suppose G is of type A . Consider the total flag variety $X = \amalg_P G/P$ (the disjoint union is taken over all parabolic subgroups). Consider the direct sum $\mathcal{O} = \oplus_{P,Q} {}^Q \mathcal{O}^P$ of double structure \mathfrak{h} -sheaves supported on the respective double moment graphs (see Example 5.5). We call \mathcal{O} the total double structure \mathfrak{h} -sheaf of the group G .

As an immediate consequence of Theorem 9.1 we have

$$\Gamma(\mathcal{O}) = {}^W\mathfrak{h}_T(X \times X) = \bigoplus_{P,Q} {}^W\mathfrak{h}_T(G/Q \times G/P),$$

therefore, we obtain

Corollary 10.5. *If G is of type A, the correspondence product induces the (non-commutative) ring structure on the space of global sections $\Gamma(\mathcal{O}) = \oplus_{P,Q} \Gamma(Q\mathcal{O}^P)$ of the total double structure \mathfrak{h} -sheaf of G .*

Moreover, the ring $\Gamma(\mathcal{O})$ is isomorphic to the endomorphism ring of the W -invariant T -equivariant motive of the total flag X , i.e., we have

$$(\Gamma(\mathcal{O}), \circ) \simeq \text{End}_{w\mathfrak{M}_T}([X]).$$

Example 10.6. Suppose $\mathfrak{h} = CH(-; \mathbb{Q})$. Then ${}^W\mathfrak{h}_T(G/Q \times G/P) \simeq S^{W_Q} \otimes_{S^W} S^{W_P}$ is a free S^W -module of rank $r(Q, P) = |{}^QW^P|$. Therefore,

$$({}^W\mathfrak{h}_T(G/Q \times G/P), \circ) \simeq M_{r(Q,P)}(S^W) \quad \text{and} \quad \Gamma(\mathcal{O}) \simeq M_r(S^W), \quad r = \sum_{(Q,P)} r(Q, P).$$

Product of global sections. Given $\phi \in {}^W\mathfrak{h}_G(G/Q \times G/P)$ and $\psi \in {}^W\mathfrak{h}_G(G/P \times G/H)$ so that ϕ and ψ are global sections of the respective double structure sheaves, we associate a new element $\psi \circ \phi \in {}^W\mathfrak{h}_G(G/Q \times G/H)$ given by the correspondence product, which is a global section of the respective double structure sheaf as well. We want to describe this product explicitly.

To do this we extend the arguments of [CNZ, §8] (were the case $P = Q$ was considered). We first investigate the Q -module $\text{Hom}_{Q_W}(Q_{W/W_Q}^*, Q_{W/W_P}^*)$. Let $\{f_v\}$ denote the standard basis of the free Q -module Q_{W/W_Q}^* or Q_{W/W_P}^* (we use the same letter f for these bases). Since the Q_W -module (via the \odot -action) Q_{W/W_Q}^* is generated by f_e (we have $w \odot f_e = f_{\bar{w}}$ for any $w \in W$), any homomorphism $\phi \in \text{Hom}_{Q_W}(Q_{W/W_Q}^*, Q_{W/W_P}^*)$ is uniquely determined by its value on f_e that is

$$\phi(f_e) = (b_v) = \sum_{v \in W^P} b_v f_v, \quad b_v \in Q.$$

Since ϕ is Q_W -linear, it has to satisfy

$$w \odot \phi(f_e) = \phi(w \odot f_e) = \phi(f_{\bar{w}}) = \phi(f_e)$$

for all $w \in W_Q$, which translates as

$$w \odot \sum_{v \in W^P} b_v f_v = \sum_{v \in W^P} w(b_v) f_{\bar{w}v} = \sum_{v \in W^P} b_v f_v \quad \text{for all } w \in W_Q.$$

Observe that the latter gives the condition (ii) of Theorem 9.1(A). In other words, (ii) guarantees that (b_v) define a Q_W -equivariant homomorphism $Q_{W/W_Q}^* \rightarrow Q_{W/W_P}^*$.

Similarly, $\psi \in {}^W\mathfrak{h}_G(G/P \times G/H)$ is also determined by its value at f_e . Let

$$\psi(f_e) = (c_u) = \sum_{u \in W^H} c_u f_u, \quad c_u \in Q,$$

hence, we have (ϕ and ψ are Q_W -equivariant)

$$\begin{aligned} \psi \circ \phi(f_e) &= \psi\left(\sum_{v \in W^P} b_v f_v\right) = \sum_{v \in W^P} b_v \psi(f_v) = \sum_{v \in W^P} b_v \psi(v \odot f_e) = \\ &= \sum_{v \in W^P} b_v (v \odot \psi(f_e)) = \sum_{v \in W^P} b_v \sum_{u \in W^H} v(c_u) f_{\bar{v}u}. \end{aligned}$$

Summarizing, we obtain the following

Proposition 10.7. *The correspondence product $(a_w) = (c_u) \circ (b_v)$ is given by*

$$\sum_{\{v \in W^P, u \in W^H \mid \overline{vu} = w\}} b_v v(c_u) = a_w, \text{ for all } w \in W^H.$$

Observe that tensoring with Q induces an embedding

$${}^W \mathbf{h}_G(G/Q \times G/P) = \mathrm{Hom}_{S_W}(\mathbb{D}_{F,Q}^*, \mathbb{D}_{F,P}^*) \hookrightarrow \mathrm{Hom}_{Q_W}(Q_{W/W_Q}^*, Q_{W/W_P}^*).$$

We now investigate its image. Recall that

$$\mathbb{D}_{F,Q}^* \subset S_{W/W_Q}^* \subset Q_{W/W_Q}^*$$

is a \mathbb{D}_F -module generated by the class of a point $[pt] = x_{\Pi/Q} f_e \in S_{W/W_Q}^*$, where $x_{\Pi/Q} = x_{\Pi}/x_Q$, $x_Q = \prod_{\alpha \in \Sigma_Q^-} x_\alpha$ and $x_{\Pi} = \prod_{\alpha \in \Sigma^-} x_\alpha$. Therefore, any ϕ is determined by its value on $[pt]$. On the other hand, $\phi([pt])$ belongs to $\mathbb{D}_{F,P}^*$ as an element of $S_{W/W_P}^* \subset Q_{W/W_P}^*$ if it satisfies the criteria (i) of Theorem 9.1(A). Combining these together, we obtain that

Lemma 10.8. *The element (global section of the localized double structure sheaf) $\phi \in \mathrm{Hom}_{Q_W}(Q_{W/W_Q}^*, Q_{W/W_P}^*)$ comes from $\mathrm{Hom}_{S_W}(\mathbb{D}_{F,Q}^*, \mathbb{D}_{F,P}^*)$ if and only if the coefficients $b_v \in Q$ satisfy conditions*

$$b'_v = x_{\Pi/Q} b_v \in S \text{ and } x_{v(\alpha)} \mid (b'_v - b'_{\overline{s_{v(\alpha)}v}}) \text{ for all } \alpha \notin \Sigma_P.$$

REFERENCES

- [BMP] T. Braden, R. MacPherson, From moment graphs to intersection cohomology, *Math. Annalen* **321** (2001), 533–551.
- [Br97] M. Brion, Equivariant Chow groups for torus actions, *Transform. Groups* **2** (1997), no.3, 225–267.
- [CNZ] B. Calmès, A. Neshitov, K. Zainoulline, Relative equivariant motives versus modules, arXiv:1609.06929.
- [CPZ] B. Calmès, V. Petrov, K. Zainoulline, Invariants, torsion indices and cohomology of complete flags, *Ann. Sci. École Norm. Sup. (4)* **46** (2013), no.3, 405–448.
- [CZZ] B. Calmès, K. Zainoulline, C. Zhong, A coproduct structure on the formal affine Demazure algebra, *Math. Zeitschrift* **282** (2016), no.3, 1191–1218.
- [CZZ1] B. Calmès, K. Zainoulline, C. Zhong, Push-pull operators on the formal affine Demazure algebra and its dual, arXiv:1312.0019.
- [CZZ2] B. Calmès, K. Zainoulline, C. Zhong, Equivariant oriented cohomology of flag varieties, *Documenta Math. Extra Volume: Alexander S. Merkurjev's 60th Birthday* (2015), 113–144.
- [CM06] V. Chernousov, A. Merkurjev, Motivic decomposition of projective homogeneous varieties and the Krull-Schmidt theorem, *Transform. Groups* **11** (2006), no.3, 371–386.
- [Cu85] C.W. Curtis, On Lusztig's isomorphism theorem for Hecke algebras, *J. Algebra* **92** (1985), 348–365.
- [EG98] D. Edidin, W. Graham, Equivariant intersection theory, *Invent. Math.* **131** (1998), no.3, 595–634.
- [Fi11] P. Fiebig, Sheaves on affine Schubert varieties, modular representations, and Lusztig's conjecture, *J. Amer. Math. Soc.* **24** (2011), 133–181.
- [Fi08] P. Fiebig, Sheaves on moment graphs and a localization of Verma flags, *Adv. Math.* **217** (2008), 683–712.
- [F15] P. Fiebig, M. Lanini, Filtered modules on moment graphs and periodic patterns, arXiv:1504.01699v1.
- [GKV] V. Ginzburg, M. Kapranov, E. Vasserot, Residue construction of Hecke algebras, *Advances in Math.* **128** (1997), no.1, 1–19.
- [Go14] R. Gonzales, Rational smoothness, cellular decompositions and GKM theory, *Geometry & Topology* **18** (2014), 291–326.

- [GKM] M. Goresky, R. Kottwitz, R. MacPherson, Equivariant cohomology, Koszul duality, and the localization theorem, *Invent. Math.* **131** (1998), 25–83.
- [GH04] V. Guillemin, T. Holm, GKM theory for torus actions with nonisolated fixed points, *Int. Math. Res. Not.* **40** (2004), 2105–2124.
- [GHZ] V. Guillemin, T. Holm, C. Zara, GKM description of the equivariant cohomology ring of a homogeneous space, *J. Algebraic Combin.* **23** (2006), no.1, 21–41.
- [GR13] N. Ganter, A. Ram, Generalized Schubert Calculus, *J. Ramanujan Math. Soc.* **28A** (Special Issue 2013), 1–42.
- [HM13] J. Heller, J. Malagón-López, Equivariant algebraic cobordism, *J. Reine Angew. Math.* **684** (2013), 87–112.
- [HMSZ] A. Hoffnung, J. Malagón-López, A. Savage, K. Zainoulline, Formal Hecke algebras and algebraic oriented cohomology theories. *Selecta Math.* **20** (2014), no.4, 1213–1245.
- [KK90] B. Kostant, S. Kumar, T -Equivariant K -theory of generalized flag varieties. *J. Differential Geometry* **32** (1990), 549–603.
- [KK86] B. Kostant, S. Kumar, The nil Hecke ring and cohomology of G/P for a Kac-Moody group G . *Advances in Math.* **62** (1986), 187–237.
- [Kr12] A. Krishna, Equivariant cobordism for torus actions. *Advances in Math.* **231** (2012), no.5, 2858–2891.
- [La12] M. Lanini, Kazhdan-Lusztig combinatorics in the moment graph setting, *J. of Algebra* **370** (2012), 152–170.
- [LZZ] C. Lenart, K. Zainoulline, C. Zhong, Parabolic Kazhdan-Lusztig basis, Schubert classes and equivariant cohomology. arXiv:1608.06554.
- [LM07] M. Levine, F. Morel, Algebraic cobordism, Springer Monographs in Math., Springer, Berlin, 2007, xii+244pp.
- [NPSZ] A. Neshitov, V. Petrov, N. Semenov, K. Zainoulline, Motivic decompositions of twisted flag varieties and representations of Hecke-type algebras, arXiv:1505.07083.
- [PS16] V. Petrov, N. Semenov, Rost motives, affine varieties, and classifying spaces. to appear in *J. of London Math. Soc.*
- [Ro03] I. Rosu, Equivariant K -theory and equivariant cohomology with an appendix by I. Rosu and A. Knutson, *Math. Zeitschrift* **243** (2003), 423–448.
- [St75] R. Steinberg, On a Theorem of Pittie. *Topology* **14** (1975), 173–177.
- [St05] J. Stembridge, Tight quotients and double quotients in the Bruhat order, *Electron. J. Combinatorics* **11** (2005), no.2, 1–41.
- [To99] B. Totaro, The Chow ring of a classifying space. Algebraic K -theory (Seattle, WA, 1997), 249–281, Proc. Sympos. Pure Math., 67, Amer. Math. Soc., Providence, RI, 1999.
- [VV03] G. Vezzosi, A. Vistoli, Higher algebraic K -theory for actions of diagonalizable algebraic groups. *Invent. Math.* **153** (2003), no.1, 1–44.
- [Vi07] A. Vistoli, On the cohomology and the Chow ring of the classifying space PGL_p , *J. Reine Angew. Math.* **610** (2007), 181–227.
- [ZZ17] G. Zhao, C. Zhong, Geometric representations of the formal affine Hecke algebra, to appear in *Advances in Math.*

(Rostislav Devyatov) DEPARTMENT OF MATHEMATICS AND STATISTICS, UNIVERSITY OF OTTAWA, 585 KING EDWARD STREET, OTTAWA, ON, K1N 6N5, CANADA

E-mail address: `deviatov@mccme.ru`

URL: <http://www.mccme.ru/~deviatov/>

(Martina Lanini) DIPARTIMENTO DI MATEMATICA, UNIVERSITÀ DI ROMA TOR VERGATA, VIA DELLA RICERCA SCIENTIFICA, 00133 ROME, ITALY

E-mail address: `lanini@mat.uniroma2.it`

URL: <https://sites.google.com/site/martinalanini5/home>

(Kirill Zainoulline) DEPARTMENT OF MATHEMATICS AND STATISTICS, UNIVERSITY OF OTTAWA, 585 KING EDWARD STREET, OTTAWA, ON, K1N 6N5, CANADA

E-mail address: `kirill@uottawa.ca`

URL: <http://mysite.science.uottawa.ca/kzaynull/>